

Secondary Kodaira-Spencer classes and nonabelian Dolbeault cohomology

Carlos Simpson
 CNRS, Laboratoire Emile Picard
 Université Toulouse III
 31062 Toulouse CEDEX, France
 carlos@picard.ups-tlse.fr

One of the nicest things about variations of Hodge structure is the “infinitesimal variation of Hodge structure” point of view [25]. A variation of Hodge structure $(V = \bigoplus V^{p,q}, \nabla)$ over a base S gives rise at any point $s \in S$ to the *Kodaira-Spencer map*

$$\kappa_s : T(S)_s \rightarrow \text{Hom}(V_s^{p,q}, V_s^{p-1,q+1}).$$

In the geometric situation of a family $X \rightarrow S$ we have

$$V_s^{p,q} = H^q(X_s, \Omega_{X_s}^p),$$

and the Kodaira-Spencer map is given by cup-product with the Kodaira-Spencer deformation class

$$T(S)_s \rightarrow H^1(X_s, T(X_s)).$$

The Kodaira-Spencer map is a component of the connection ∇ . In particular, this implies that if $\kappa_s \neq 0$ then the connection ∇ is nontrivial with respect to the Hodge decomposition. Various Hodge-theory facts imply that the global monodromy must be nontrivial in this case. We can be a bit more precise: if $u \in V^{p,q}$ is a vector such that $\kappa_s(v)(u) \neq 0$ for some tangent vector $v \in T(S)_s$, then u cannot be preserved by the global monodromy. Thus a local calculation (which actually only depends on the first-order deformation of X_s) implies a global fact. In particular this global fact would hold for any family of varieties X' over any base S' , such that the new family osculates to order 1 with the original one (say as a map from S' into the moduli stack of the fibers). A particularly nice aspect of this situation is that the Kodaira-Spencer map is defined on the Dolbeault cohomology $H^q(X_s, \Omega^p)$ and in particular it is obtained involving only algebraic-geometric calculations (just a cup-product with the deformation class)—no analytic considerations are needed.

The goal of this paper is to calculate an example showing a similar type of behavior with a secondary Kodaira-Spencer class coming from nonabelian cohomology with coefficients in the complexified 2-sphere $T = S^2 \otimes \mathbf{C}$. For such a T (or any other coefficient stack similar in nature) we define the *nonabelian Dolbeault cohomology of X with coefficients in T* , denoted $\text{Hom}(X_{\text{Dol}}, T)$. When X varies in a family parametrized by a base

S then we show how to define a secondary class which is a map

$$\bigwedge^2 T(S)_s \rightarrow \pi_1(\text{Hom}(X_{Dol}, T)) = H_{Dol}^2(X)/(\eta)$$

where η is the class in $H_{Dol}^2(X)$ pulled back from the tautological class on T by the map $X_{Dol} \rightarrow T$ which we take as basepoint. The secondary class is defined when the primary Kodaira-Spencer classes vanish.

It seems likely, although we don't show that here, that our secondary class is just a quadruple Massey product

$$\alpha \wedge \beta \mapsto \{\eta, \eta, \alpha, \beta\}$$

for $\alpha, \beta \in H^1(X, TX)$ and $\eta \in H_{Dol}^2(X)$ with $\eta \cup \eta = 0$. Instead of interpreting the secondary class this way, we calculate it directly using Mayer-Vietoris arguments for nonabelian cohomology and reducing to calculations in abelian cohomology.

Our main purpose in the present paper is to calculate the secondary class in a specific, somewhat instructive, example. We look at the family of varieties X obtained by blowing up a point $P \in Z$ on a smooth surface Z . This family is parametrized in an obvious way by Z . If Z is simply connected, then any standard Hodge-theoretic information related to X must be independent of the basepoint because it would vary in a variation of (mixed) Hodge structure parametrized by Z , and such a variation is forcibly constant. However, we will show that the secondary Kodaira-Spencer class for nonabelian Dolbeault cohomology is nonzero, if Z has a nonzero holomorphic 2-form.

Before getting to the example, we will discuss some general aspects of nonabelian cohomology with emphasis on the case of Dolbeault cohomology. We start with a quick review of n -stacks in §1. Then we define the notion of “connected very presentable shape” in §2 by looking at maps to n -stacks T which are 0-connected, with $\pi_1(T)$ an affine algebraic group scheme, and $\pi_i(T)$ vector spaces, for $i \geq 2$. The next step in §3 is to set X_{Dol} equal to the 1-stack over X whose fiber is $K(\widehat{TX}, 1)$ where $\widehat{TX}$ is the completion of the tangent bundle of X along the zero-section. We define the *nonabelian Dolbeault cohomology of X with coefficients in an n -stack T* to be $\text{Hom}(X_{Dol}, T)$. We look at this particularly for connected very presentable T . This gives rise to the *Dolbeault shape* which is the $(n+1)$ -functor

$$T \mapsto \text{Hom}(X_{Dol}, T)$$

on connected very presentable T . Sections 4 and 5 treat group actions and secondary classes in the n -stack situation. In §5 we define the secondary Kodaira-Spencer class in general. Then in §6 we look at a particular case of T , namely the *complexified 2-sphere*, defined by the conditions that $\pi_2 = \pi_3 = \mathcal{O}$ and having nontrivial Whitehead product.

With these preliminary steps done, we get to our example in §§7-8. The main result is Theorem 7.1. It says that if X is the blowup of a surface Z at a point P then the

secondary Kodaira-Spencer class for nonabelian Dolbeault cohomology $Hom(X_{Dol}, T)$ with coefficients in the complexified 2-sphere T , is Serre dual to the evaluation map of holomorphic 2-forms at P . In particular if $H^0(Z, \Omega_Z^2) \neq 0$ then this class is nonzero.

At the end in two appendices, we will discuss some topics from [39] but in greater detail. First, the “Breen calculations” giving the cohomology of $K(\mathcal{O}, n)$ or more generally $K(V/S, n)$ for a vector sheaf V over a base scheme S . Then we discuss representability of certain shapes. These topics come up in a few places in the body of the paper, which is the reason for the appendices; for an introduction we refer the reader to the appendices.

Without going into great bibliographic detail, we point out here some recent papers which seem to be somewhat related. Karpishpan [29] [30] treats higher-order Kodaira-Spencer mappings, i.e. the higher order derivatives of the period map. The same is treated by Ran [34] and Esnault-Viehweg [17]. Biswas defines secondary invariants for families of Higgs bundles [6]. Also Bloch and Esnault treat algebraic Cherns-Simons classes [7], which are types of secondary classes.

I would like to thank Mark Green for an inspiring question—albeit one which the present paper doesn’t answer. He asked whether there are examples of families of varieties where the variation of Hodge structure on the cohomology is constant, but where the variation of mixed Hodge structure on the homotopy groups is nontrivial. In the absence of an answer to that question (which is very interesting), the typical mathematician’s reply is to change the question—in this case, to look for an example where even the mixed Hodge structures on the homotopy groups remain constant, but where the Hodge filtration on the full homotopy type is nonconstant.¹

Notation

We always work in characteristic 0. In order to simplify notation we use $\mathbf{C}$ as the ground field (i.e. $Spec(\mathbf{C})$ as base scheme), but everything we say would work equally well over any ground field of characteristic 0. Let $Sch/\mathbf{C}$ denote the site of schemes of finite type over $Spec(\mathbf{C})$ with the etale topology.

The structure sheaf $\mathcal{O}$ on $Sch/\mathbf{C}$ is the sheaf defined by

$$\mathcal{O}(Y) := \Gamma(Y, \mathcal{O}_Y).$$

It is represented by the affine line $\mathbf{A}^1$, in other words it is represented by the 1-dimensional vector space $\mathbf{C}$. A finite dimensional vector space represents a sheaf of the form $\mathcal{O}^a$.

¹ After the first version of this paper, Richard Hain pointed out to me the following example answering Mark Green’s original question. The example comes from a paper of Carlson, Clemens, and Morgan ([14], p. 330). Let $C \subset \mathbf{P}^3$ be an embedded curve of positive genus. For points $p, q \in C$, let $X_{p,q}$ be the 3-fold obtained by first blowing up p and q and then blowing up the strict transform of C . The family of $X_{p,q}$ parametrized by (an open subset of) $C \times C$ has constant variation of Hodge structures on the cohomology but, according to [14] the variation of MHS on the homotopy is nonconstant.

1. Basic remarks about n -stacks

We make some brief remarks about n -stacks as we shall use them in this paper. For all details the reader is referred to the following references:

- for the history and basic notions of simplicial presheaves: Brown [12], Illusie [24], Jardine [27];
- for the history and basic notions of 1-stacks: Artin [2], Deligne-Mumford [16], and Laumon-Moret-Bailly [31];
- for cohomological theory using simplicial presheaves: Thomason [47];
- for a “homotopy coherent” approach: Cordier-Porter [15] and also [38];
- for n -categories and n -stacks: Grothendieck [20], Breen [10], Gordon-Power-Street [19], Tamsamani [46], Baez-Dolan [3], and several papers of the author.

The first main remark is that we shall almost always be concerned with n -stacks of n -groupoids, and following the intuition put forth in [20], an n -groupoid is the same thing (up to homotopy) as a topological space whose homotopy groups vanish in degrees $i > n$ —we call such a space *n -truncated*. Thus it is safe to replace n -groupoids everywhere by n -truncated topological spaces or, again equivalently, n -truncated simplicial sets. In this point of view, an n -stack (on the site $Sch/\mathbf{C}$ which is fixed throughout) is just a presheaf of simplicial sets otherwise known as a *simplicial presheaf*, which is object-by-object n -truncated. Once one has made the passage to simplicial presheaves, the truncation condition is no longer crucial (although it often facilitates arguments and many things in the literature are only stated in this case or under a complementary hypothesis about vanishing cohomological dimension). Thus, when we speak of “ n -stacks”, one way to read this is in terms of the homotopy theory of simplicial presheaves, see [12] [24] [27] [47].

Another reading would plunge directly into the theory of n -categories and n -stacks, imposing the groupoid condition along the way. For $n \leq 3$ (which in the end is the case we treat in the present paper) this can be had in a relatively formulaic way in [10] and [19]. For arbitrary n , see [46] [43], but unfortunately the n -stack part of this theory still needs to be worked out a bit more.

The only place where we make reference to n -categories which are not n -groupoids is when we look at the $n + 1$ -category $nSTACK$ of n -stacks (of groupoids). This $n + 1$ -category has the property of being *1-groupic*, i.e. the morphism n -categories are n -groupoids. One can safely replace the morphism n -groupoids by spaces or simplicial sets, and one obtains the notion of *Segal category* [45], motivated by Segal’s delooping machine [35]. This notion came into higher category theory in Tamsamani’s definition of n -category [46]. Thus $nSTACK$ may be considered as a Segal category. This fits in relatively nicely with the simplicial presheaf point of view; in fact this Segal category comes from the simplicial category of fibrant and cofibrant objects in Jardine’s closed

model category of simplicial presheaves. This comes up in looking at the functoriality in T of the construction $Hom(X_{Dol}, T)$. In one other place we refer to the $n + 1$ -stack $nSTACK$ of n -stacks, which is discussed somewhat in [43] and [44]; we don't get any further into the general theory here.

Currently, the “simplicial presheaves” alternative is the most accessible (it is also historically the first, dating from [12]). In terms of simplicial presheaves, an n -stack is a simplicial presheaf on $Sch/\mathbf{C}$ which is object-by-object n -truncated. If X, Y are simplicial presheaves, then we obtain a simplicial presheaf $Hom(X, Y)$ by first replacing Y by a fibrant object [27], then looking at the internal Hom of simplicial presheaves. In particular this gives a simplicial set $Hom(X, Y)(Spec(\mathbf{C}))$ and this family of simplicial sets makes the simplicial presheaves into a simplicial category (or Segal category) which we denote $nSTACK$. When we speak of morphisms between n -stacks, the above procedure is always understood, i.e. we always replace the target by a fibrant object.

Given a presheaf of n -groupoids or presheaf of spaces, we often want to take the “associated n -stack”. What this means depends somewhat on the point of view which is taken. If one works with objects in a closed model category such as that of simplicial presheaves, then this just means to consider the object as an element of the closed model category. One might also want to say that it means to replace the object by a weakly equivalent fibrant object. Finally there is an intermediate notion based on enforcing the global descent condition but not the local fibrant condition. It doesn't really matter which point of view we adopt, since when looking at morphisms to a given object, we always replace it by an equivalent fibrant object anyway.

The “yoga” of the situation is that one can do topology with n -stacks instead of spaces (or more precisely n -truncated spaces). In particular, all standard constructions and results in algebraic topology carry over to n -stacks. Most of these are contained somewhere in the literature referred to above; but if not, we don't give proofs here as that would get beyond the scope of the present paper.

There is basically only one slight “twist” which is not present in the topological case: this is that the 0-truncated objects can be topologically nontrivial, i.e. can have cohomology. In the usual topological case, the 0-truncated objects are just the disjoint unions of contractible components and these make no significant contribution to homotopy. In the case of n -stacks over a site, one can have 0-stacks, i.e. sheaves of sets, with nontrivial cohomology (this is the case of a smooth projective variety X , for example); and similarly there are sheaves of groups over $*$ (the site itself) which can have nontrivial cohomology. The upshot of all this is that when it comes to choosing basepoints for an n -stack T , one must choose first an object $Y \in Sch/\mathbf{C}$ and then choose a basepoint $t \in T(Y)$. (In other words, if we look only at basepoints in $T(Spec(\mathbf{C}))$ we might be missing some topology.)

The first place where the previous paragraph has an impact is in the notion of *homotopy groups*. If T is an n -stack then for any $Y \in Sch/\mathbf{C}$ and $t \in T(Y)$ we obtain a presheaf,

denoted in utmost precision by

$$\pi_i^{\text{pre}}(T|_{Sch/Y}, t)$$

but which we often shorten to $\pi_i^{\text{pre}}(T, t)$. This is a presheaf of groups over the site Sch/Y , abelian if $i \geq 2$. On the other hand the presheaf $\pi_0^{\text{pre}}(T)$ is defined absolutely as a presheaf of sets over $Sch/\mathbf{C}$.

The definition which is fundamental to the theory is that we define

$$\pi_i(T|_{Sch/Y}, t)$$

to be the sheaf associated to the presheaf $\pi_i^{\text{pre}}(T|_{Sch/Y}, t)$. Similarly $\pi_0(T)$ is the sheaf of sets associated to the presheaf $\pi_0^{\text{pre}}(T)$. These, and not the presheaf versions, are the only thing we care about. This is formalized by saying that a morphism $f : T \rightarrow T'$ is called a *weak equivalence* [24] if for all $Y \in Sch/\mathbf{C}$ and $t \in T(Y)$, the resulting morphisms

$$\pi_i(T|_{Sch/Y}, t) \rightarrow \pi_i(T'|_{Sch/Y}, f(t))$$

are isomorphisms of sheaves on Sch/Y (resp. $\pi_0(T) \rightarrow \pi_0(T')$ is an isomorphism of sheaves of sets on $Sch/\mathbf{C}$). The theory is localized by this notion of equivalence, in other words T and T' are thought of as equivalent if there is a weak equivalence between them. Jardine constructs a closed model category which takes this into account [27]. This leads, in particular, to the right notion of morphism, namely we only look at morphisms whose target is a fibrant object; if necessary, a target object is replaced by a weakly equivalent fibrant object. Without further mentioning this, we make the convention that whenever we speak of morphisms between n -stacks, the target object is made fibrant.

For a general site, one can have a connected stack T (i.e. $\pi_0(T) = *$) but where the global section space of T is empty, or nonconnected. However, in the present case we are working in the etale topology over an algebraically closed field $\mathbf{C}$. In this case we have the implication

$$\pi_0(T) = * \Rightarrow \pi_0(T(Spec(\mathbf{C}))) = *.$$

Indeed, the etale coverings of $Spec(\mathbf{C})$ are trivial, so there is no change over the object $Spec(\mathbf{C})$ when one passes from the presheaf $\pi_0^{\text{pre}}(T)$ to the associated sheaf.

If T is connected, then, we can choose a basepoint $t \in T(Spec(\mathbf{C}))$ which is unique up to homotopy, so the sheaf of groups $\pi_1(T, t)$ is uniquely defined up to global conjugacy. If Y is any scheme and $t' \in T(Y)$ then locally on Y , t' is equivalent to $t|_Y$ so $\pi_1(T, t')$ is locally over Y equivalent to the restriction of $\pi_1(T, t)$ to Y . Thus in this case, the fundamental group sheaf $\pi_1(T, t)$ over $Sch/\mathbf{C}$ gives a relatively accurate picture of the 1-type of T .

In particular, it makes sense to require that T be 1-connected, that is that $\pi_0(T) = *$ and $\pi_1(T, t) = \{1\}$ for the basepoint $t \in T(Spec(\mathbf{C}))$. If T is 1-connected then for any

scheme Y and $t' \in T(Y)$, $\pi_1(T, t') = \{1\}$. Furthermore, in this case the fundamental group of $T(\text{Spec}(\mathbf{C}))$ is trivial (the cohomological contributions from the higher homotopy vanish because etale cohomology of $\text{Spec}(\mathbf{C})$ is trivial). Therefore the basepoint $t \in T(\text{Spec}(\mathbf{C}))$ is well-defined up to unique homotopy.

We now describe the standard topological constructions which we shall use for n -stacks. The first is the notion of *homotopy fiber product*. If $A \rightarrow B \leftarrow C$ are morphisms of n -stacks then we obtain the *homotopy fiber product* $A \times_B C$ with a diagram

$$\begin{array}{ccc} A \times_B C & \rightarrow & A \\ \downarrow & & \downarrow \\ C & \rightarrow & B \end{array}$$

together with a homotopy of commutativity of the diagram. These data are essentially well-defined (in the sense that they are well-defined up to homotopy which is itself well-defined up to homotopy $\dots$). In the simplicial presheaf theory, the homotopy fiber product is obtained by replacing one of the two morphisms by a fibrant morphism and then taking the usual fiber product. In the n -category theory, see [44].

Suppose $f : A \rightarrow B$ is a morphism and suppose $b \in B(\text{Spec}(\mathbf{C}))$. We can think of b as a morphism $b : * \rightarrow B$ where $*$ denotes the constant presheaf with values the 1-point topological space. Define the *fiber of f over b* to be the homotopy fiber product

$$\text{Fib}(f, b) := * \times_B A.$$

If the base B is 0-connected, then as mentioned above, the choice of basepoint b exists and is unique up to a global homotopy (i.e. a path in $B(\text{Spec}(\mathbf{C}))$). Thus we can denote by $\text{Fib}(f)$ the fiber over this b , bearing in mind that it is defined up to the conjugation action of $\pi_1(B(\text{Spec}(\mathbf{C})))$. If B is 1-connected the choice of basepoint b is unique up to unique homotopy, so $\text{Fib}(f)$ is well-defined up to homotopy. We call this the *homotopy fiber of f* .

We say that

$$A \rightarrow B \rightarrow C$$

is a *fiber sequence* if C is 1-connected or if we are otherwise given a basepoint $c \in C(\text{Spec}(\mathbf{C}))$, if we are given a homotopy between the composition $A \rightarrow C$ and the constant map at the basepoint c , and if the map $A \rightarrow B$ together with this homotopy induce an equivalence between A and $\text{Fib}(B \rightarrow C, c)$.

A morphism $T \rightarrow R$ is said to be a *locally constant fibration with fiber F* if for every scheme $Y \rightarrow R$, locally on Y (in the etale topology) we have $Y \times_R T \cong Y \times F$. In the usual topological case with connected base, this is vacuous. In the case of n -stacks over a site, we still have that if R is 0-connected then any morphism $T \rightarrow R$ is a locally constant fibration. However, we are often interested in cases where R is not connected (i.e. $\pi_0(R)$

is some nontrivial sheaf of sets). In these cases, being locally constant condition is a nontrivial additional condition.

The next general type of operation we discuss is *truncation*. If T is an n -stack and if $m \leq n$ then we obtain an m -stack $\tau_{\leq m}T$ (i.e. a simplicial presheaf which is m -truncated, in other words has homotopy group sheaves vanishing in degrees $> m$) together with a morphism of n -stacks

$$T \rightarrow \tau_{\leq m}T$$

which induces an isomorphism on homotopy group sheaves in degrees $i \leq m$. The m -stack $\tau_{\leq m}T$ together with this morphism are essentially well-defined. We can construct $\tau_{\leq m}T$ as the m -stack associated to the presheaf of spaces

$$(\tau_{\leq m}^{\text{pre}}T)(Y) := \tau_{\leq m}(T(Y))$$

where the truncation on the left is just truncation of topological spaces (also known as the coskeleton operation).

A first example of truncation is the sheaf of sets $\pi_0(T) = \tau_{\leq 0}T$.

If $T \rightarrow R$ is a morphism of n -stacks then there is a relative (or “fiberwise”) version of the truncation denoted $\tau_{\leq m/R}(T) \rightarrow R$. This is defined by the property that for any scheme Y and morphism $Y \rightarrow R$,

$$\tau_{\leq m/R}(T) \times_R Y = \tau_{\leq m}(T \times_R Y).$$

Using the operations of truncation and homotopy fiber products, we obtain the *Postnikov tower*. If T is an n -stack then we have morphisms

$$\begin{aligned} T &\rightarrow \dots \rightarrow \tau_{\leq m}T \\ &\rightarrow \tau_{\leq m-1}T \rightarrow \dots \rightarrow \pi_0(T). \end{aligned}$$

In order to describe the stages in this tower of maps, we need a few more notions.

Suppose Y is a scheme and L is a sheaf of groups over Y . Fix $m \leq n$ and suppose L is abelian if $m \geq 1$. Then we can construct the simplicial presheaf $K^{\text{pre}}(L, m)$ on Sch/Y by the standard construction applied to L ; let $K(L, m)$ be the associated stack. Note that $K(L, m)$ has a chosen basepoint section (over Y) which we denote by 0 , and $\pi_i(K(L, m), 0) = 0$ for $i \neq m$, and it is $= L$ for $i = m$. Furthermore these properties characterize $K(L, m)$ essentially uniquely.

The $K(L, m)$ on the site Sch/Y corresponds to an n -stack on $Sch/\mathbf{C}$ with morphism to Y (where Y is considered as a 0-stack or sheaf of sets), which we denote by $K(L/Y, m) \rightarrow Y$.

We can do the same construction relative to any n -stack but for this we need to have a notion corresponding to sheaf of (abelian) groups. If A is an n -stack then a *local system*

of (abelian) groups) on A is a morphism $L \rightarrow A$ with relative group structure, which is relatively 0-truncated (i.e. for any scheme Y and map $Y \rightarrow A$, the homotopy fiber product $Y \times_A L$ is 0-truncated). This is equivalent to the data for every Y of a local system of (abelian) groups L_Y over $A(Y)$, together with restriction morphisms $L_Y|_{A_{Y'}} \rightarrow L_{Y'}$ for $Y' \rightarrow Y$, satisfying the obvious associativity condition.

Caution: if X is a sheaf of sets represented by a scheme, then a local system over X (according to the above terminology) is the same thing as a sheaf of (abelian) groups over X . It doesn't have anything to do with the notion of "flat vector bundle" over X .

If $L \rightarrow A$ is a local system of abelian groups then we obtain a morphism

$$K(L/A, n) \rightarrow A,$$

whose homotopy fiber over any $a \in A(Y)$ is the Eilenberg-MacLane n -stack $K(L|_Y, n)$ over Y . For $n = 1$ we can make do with any local system of groups not necessarily abelian.

There is a standard fibration sequence relative to A

$$K(L/A, m) \rightarrow A \rightarrow K(L/A, m+1),$$

in other words

$$K(L/A, m) = A \times_{K(L/A, m+1)} A.$$

Using this we obtain the usual description of the stages in the Postnikov tower: if $m \geq 2$ then, setting $A := \tau_{\leq m-1}T$ there is a local system L of abelian groups over A and a section $ob : A \rightarrow K(L/A, m+1)$ such that the morphism in the Postnikov tower

$$\tau_{\leq m}T \rightarrow \tau_{\leq m-1}T = A$$

is equivalent to

$$A \times_{K(L/A, m+1)} A \rightarrow A$$

where the first morphism in the fiber product is ob and the second is 0.

The description of the first stage $\tau_{\leq 1}T \rightarrow \pi_0(T)$ is much more complicated and is basically the subject of Giraud's book [18].

Using the notion of local system we can define a relative version of the homotopy group sheaves. If $T \rightarrow R$ is a morphism of n -stacks and $s : R \rightarrow T$ is a section then we obtain local systems of groups $\pi_i(T/R, s)$ over R .

Suppose $X \rightarrow Z$ and $Y \rightarrow Z$ are morphisms of n -stacks. Then we obtain a *relative internal Hom* which is an n -stack with morphism to S ,

$$Hom(X/Z, Y/Z) \rightarrow S.$$

It is defined by the universal property that maps $A \rightarrow Hom(X/Z, Y/Z)$ are the same (in an essentially well-defined way) as maps $X \times_Z A \rightarrow Y$ over Z . For existence, if the proof

isn't contained somewhere in the literature then one might have to apply the techniques of [44]. If $Z = *$ then we get back to the usual internal $Hom(X, Y)$.

Similarly if $Y \rightarrow X \rightarrow Z$ then we obtain the *relative section stack*

$$\Gamma(X/Z, Y)$$

which is defined to be the fiber product

$$Hom(X/Z, Y/Z) \times_{Hom(Y/Z, Y/Z)} Z$$

where the second map in the fiber product is that corresponding to the identity of Y , and the first map is induced by $Y \rightarrow X$. Again if $Z = *$ we denote this simply by $\Gamma(X, Y)$.

We now come to one of the main types of observations, namely the relationship between the above objects and cohomology. See for example Thomason [47] for much of this. If A is an n -stack and L a local system of abelian groups over A then we define

$$H^i(A, L) := \pi_0 \Gamma(A, K(L/A, i)).$$

It is a sheaf of abelian groups on the site $Sch/\mathbf{C}$. Similarly if Z is an n -stack, $p : A \rightarrow Z$ a morphism and L a local system of abelian groups over A then we define

$$H^i(A/Z, L) := \tau_{\leq 0/Z} \Gamma(A/Z, K(L/A, i))$$

where $\tau_{\leq 0/Z}$ is the relative version of the truncation operation for n -stacks over Z . Note that $H^i(A/Z, L)$ is a local system of abelian groups on Z . We can also denote it by $R^i p_*(L)$.

One has the result that the cohomology defined above coincides with sheaf cohomology over simplicial objects (representing A by a simplicial object in the topos of $Sch/\mathbf{C}$). See [47] or [38]. In particular the notation $R^i p_*(L)$ coincides with the usual meaning (particularly when we are looking at A and Z which are represented by schemes, for example).

We have the formulae

$$\pi_i(\Gamma(A, K(L/A, m)), 0) = H^{m-i}(A, L)$$

and (using the relative version of homotopy groups)

$$\pi_i(\Gamma(A/Z, K(L/A, m))/Z, 0) = H^{m-i}(A/Z, L).$$

The usual results concerning cohomology of topological spaces hold for cohomology as defined above. In particular, we have cup-products, corresponding to the following

operations on Eilenberg-MacLane spaces. If L , L' and L'' are local systems of abelian groups over A and if

$$L \times_A L' \rightarrow L''$$

is a bilinear morphism (of relative abelian group objects) then we obtain morphisms

$$K(L/A, i) \times K(L'/A, j) \rightarrow K(L''/A, i+j).$$

These give cup-products in cohomology which are bilinear morphisms

$$H^i(A, L) \times H^j(A, L') \rightarrow H^{i+j}(A, L'').$$

We also have a Künneth formula. The case which we use in the present paper is as follows. Suppose X and Y are n -stacks. Then

$$H^m(X \times Y, \mathcal{O}) = \bigoplus_{i+j=m} H^i(X, H^j(Y, \mathcal{O})).$$

If $H^j(Y, \mathcal{O})$ are represented by finite dimensional vector spaces then we can write the more usual formula

$$H^m(X \times Y, \mathcal{O}) = \bigoplus_{i+j=m} H^i(X, \mathcal{O}) \otimes_{\mathcal{O}} H^j(Y, \mathcal{O}).$$

This extends to the relative case of morphisms $X \rightarrow S$ and $Y \rightarrow S$ if these families are locally trivial over the étale topology of S .

Finally, we have a Leray-Serre spectral sequence. See [47] for one way to set this up. If $f : X \rightarrow Y$ is a morphism of n -stacks and if L is a local system of abelian groups on X then we obtain a “complex” $Rf_*(L)$ on Y and the cohomology of X is the “hypercohomology” of this complex. These terms are put in quotations because one should actually interpret the notion of complex as being a fibration in spectra over Y (the raw notion of complex of local systems is not adapted to the higher homotopy involved if Y is not 0-truncated and locally cohomologically trivial). In any case we get the cohomology objects of the direct image, which are the relative homotopy group sheaves of the fibration of spectra, denoted $R^i f_*(L)$. These are local systems over Y . We have the Leray-Serre spectral sequence (cf Thomason [47])

$$E_2^{i,j} = H^i(Y, R^j f_*(L)) \Rightarrow H^{i+j}(X, L).$$

One way of approaching all of the above details is to replace any n -stack i.e. simplicial presheaf (particularly those coming in as the base in questions about cohomology) by a simplicial object whose stages are formal, possibly infinite, disjoint unions of schemes (this technique was pointed out to me by C. Teleman). Similarly, we can replace morphisms of n -stacks by morphisms of such objects. Then questions about cohomology become just

questions about cohomology of simplicial schemes with obvious modifications made to allow for the infinite number of components.

The n -stack $K(\mathcal{O}, n)$ has an infinite loop-space structure or E_∞ -structure, in other words it has an infinite delooping (the m -fold delooping is just $K(\mathcal{O}, m+n)$). This structure is the homotopical analogue of an abelian group structure: it contains a homotopy class of maps

$$K(\mathcal{O}, n) \times K(\mathcal{O}, n) \rightarrow K(\mathcal{O}, n)$$

but also higher homotopies of associativity, commutativity etc. We often think of $K(\mathcal{O}, n)$ as a homotopical group object and speak of things such as “principal bundles” for it.

The infinite loop-space structure is inherited by $\text{Hom}(\mathcal{F}, K(\mathcal{O}, n))$ for any sheaf $\mathcal{F}$.

In the first version of the paper, it was stated that the infinite loop-space structure provides a functorial decomposition into products of Eilenberg-MacLane stacks. This is not true in general. For one thing, such a decomposition may not exist, and for another it is never completely functorial. For existence, the obstruction is in the *Ext* between the various homotopy group sheaves. If such a decomposition exists, then it can be functorial up to one homotopy, but this homotopy itself will not be uniquely determined up to a second homotopy.

The statement which we actually need for the calculation in §8 below, does work, and we state it as a proposition.

Proposition 1.1 *(A) Suppose (Y, y) is a basepointed n -stack with infinite loop structure. Suppose that $\pi_i(Y, y)$ are represented by finite-dimensional vector spaces. Then there exists an equivalence*

$$\varepsilon : Y \cong Y_0 \times \dots \times Y_n$$

where

$$Y_i = K(\mathcal{O}^{a_i}, i) = K(\pi_i(Y, y), i).$$

(B) Suppose $f : Y \rightarrow Y'$ is a map of basepointed n -stacks both as in (A), and suppose f is compatible with the infinite loop structure. Let

$$\varepsilon : Y \cong Y_0 \times \dots \times Y_n, \quad \varepsilon' : Y' \cong Y'_0 \times \dots \times Y'_n$$

denote the maps given by (A) (chosen independantly of f). Then there is a homotopy making the square

$$\begin{array}{ccc} Y & \cong & Y_0 \times \dots \times Y_n \\ \downarrow & & \downarrow \\ Y' & \cong & Y'_0 \times \dots \times Y'_n \end{array}$$

commute, where the vertical arrow on the right is a product of the morphisms of Eilenberg-MacLane stacks $Y_i \rightarrow Y'_i$ induced by $f_* : \pi_i(Y, y) \rightarrow \pi_i(Y', y')$.

Proof: The short way of saying this is that the $Ext^j(V, W)$ vanish for $j > 0$ for finite dimensional vector spaces, so the obstruction to splitting vanishes. We give the following more concrete argument (which is in a certain sense just repeating the argument which will be given in 9.6 below for the vanishing of the Ext^j).

For $N > n$ we are given an $N - 1$ -connected pointed $n + N$ -stack (Z, z) with $(Y, y) = \Omega^N(Z, z)$ (this is the *ad hoc* definition of “infinite loop structure” which we use). In part (B) the map f comes from a map $g : Z \rightarrow Z'$. Thus for part (A) it suffices to obtain a decomposition

$$Z \cong Z_0 \times \dots \times Z_n$$

with

$$Z_i = K(V_i, N + i), \quad V_i := \pi_i(Y, y) = \pi_{N+i}(Z, z),$$

and for part (B) it suffices to obtain the homotopy of functoriality on the level of the morphism g .

Calculate the cohomology of Z with coefficients in $\mathcal{O}$. Using Leray-Serre spectral sequences for the stages in the Postnikov tower, and using the Breen calculations 9.2 in view of the hypothesis that the π_j are finite dimensional vector spaces, we find that for $i \leq N + n$,

$$H^j(Z, \mathcal{O}) = \text{Hom}(\pi_j(Z, z), \mathcal{O}) = V_{N-j}^*.$$

Thus we have tautological classes

$$\varepsilon_i \in H^{N+i}(Z, \pi_{N+i}(Z, z)) = H^{N+i}(Z, V_i)$$

which together provide us with a map

$$\varepsilon = (\varepsilon_0, \dots, \varepsilon_n) : Z \rightarrow K(V_0, N) \times \dots \times K(V_n, N + n).$$

This map induces an isomorphism on homotopy groups (in degrees up to $N + n$). Thus it provides the required splitting for (A).

For part (B) suppose we have a map

$$g : Z = Z_0 \times \dots \times Z_n \rightarrow Z'_0 \times \dots \times Z'_n$$

with $Z_i = K(V_i, N + i)$ and $Z'_i = K(W_i, N + i)$. Such a map corresponds, up to homotopy, to a collection of classes in $H^{N+i}(Z, W_i)$. From the Künneth formula (which can be seen by a collection of Leray-Serre spectral sequences for the projections onto the factors) we have

$$H^{N+i}(Z, W_i) = H^{N+i}(Z_i, W_i).$$

The other factors vanish again using the Breen calculations 9.2 from the fact that V_j and W_i are represented by finite dimensional vector spaces. Our map g is therefore homotopic

to a map given by the classes in $H^{N+i}(Z_i, W_i)$, i.e. a map compatible with the product decomposition. $\square$

The homotopy in part (B) is not unique: it can be changed by a map $Y \rightarrow \Omega Y'$ in other words by a collection of morphisms $Y_i \rightarrow Y'_{i+1}$.

Suppose that $\mathcal{F}$ is an n -stack such that the $H^i(\mathcal{F}, \mathcal{O})$ are represented by finite dimensional vector spaces. Then we can apply the above proposition to

$$Y := Hom(\mathcal{F}, K(\mathcal{O}, n)).$$

The decomposition of $Hom(\mathcal{F}, K(\mathcal{O}, n))$ into a product of Eilenberg-MacLane spaces

$$Hom(\mathcal{F}, K(\mathcal{O}, n)) = \prod_i K(H^{n-i}(\mathcal{F}, \mathcal{O}), i),$$

is related to the Künneth formula. A morphism $Z \rightarrow Hom(\mathcal{F}, K(\mathcal{O}, n))$ corresponds (by the definition of internal Hom) to a morphism $Z \times \mathcal{F} \rightarrow K(\mathcal{O}, n)$, in other words to a class $f \in H^n(Z \times \mathcal{F}, \mathcal{O})$. By the above product structure this class decomposes into a collection of classes $f_i \in H^i(Z, H^{n-i}(\mathcal{F}, \mathcal{O}))$. The f_i are the Künneth components of f .

* * *

For the remainder of the paper, we look at n -stacks of groupoids on $Sch/\mathbb{C}$ and unless specified otherwise, the reader may fix any $n \geq 3$ (for the calculation it suffices to take $n = 3$.)

2. Connected very presentable shape

We isolate some special n -stacks T and then use them to measure the “shape” of an arbitrary n -stack $\mathcal{F}$. In other words, take a sub- $n + 1$ -category $\mathcal{P} \subset nSTACK$ of the $n + 1$ -category of n -stacks (of groupoids, say), and look at the $n + 1$ -functor

$$\mathcal{P} \rightarrow nSTACK$$

$$T \mapsto Hom(\mathcal{F}, T).$$

We call this $n + 1$ -functor the *shape of $\mathcal{F}$ as measured by $\mathcal{P}$* . There are many possible ways to choose $\mathcal{P}$. Some reasonable parameters are to require that $\mathcal{P} \subset nSTACK$ be a full sub- $n + 1$ -category (in other words that we make no limitation on the morphisms of $\mathcal{P}$); and that the condition $T \in \mathcal{P}$ should be measured only by looking at the homotopy group sheaves $\pi_i(T, t)$. For our present purposes we start by requiring that T be connected, i.e.

$\pi_0(T) = *$. Among other things, this insures that the isomorphism classes of the higher homotopy group sheaves $\pi_i(T, t)$ be well defined.

Before describing our choice of conditions for the $\pi_i(T, t)$ we take note of the following: any $T \in \mathcal{P}$ will decompose in a Postnikov tower whose stages are $K(\pi_i, i)$. Morphisms $\mathcal{F} \rightarrow K(\pi_i, i)$ are classified by $H^i(\mathcal{F}, \pi_i)$ and more generally, one has obstruction theory for classifying the morphisms $\mathcal{F} \rightarrow T$ going up in the Postnikov tower; the obstruction classes are in $H^{i+1}(\mathcal{F}, \pi_i)$. Thus, one should choose the class of possible π_i to be a class of sheaves such that, for the $\mathcal{F}$ we are interested in, the cohomology $H^j(\mathcal{F}, \pi_i)$ has reasonable properties.

For the topic of Dolbeault cohomology, we already know how to take nonabelian H^1 with coefficients in an affine algebraic group [37] [36], and we know how to take higher Dolbeault cohomology with coefficients in $\mathbf{C}$ or more generally in a finite-dimensional complex vector space. This suggests that our conditions should be that π_1 be an affine algebraic group, and π_i be represented by finite-dimensional vector spaces for $i \geq 2$.

Recall from [41] and [42] that a *connected very presentable n -stack* T is an n -stack of groupoids T on $Sch/\mathbf{C}$ subject to the following conditions:

(connectedness): $\pi_0(T) = *$ as a sheaf of sets on $Sch/\mathbf{C}$;

(very presentability): if $t \in T(Spec(\mathbf{C}))$ is a basepoint (which we assume exists) then $\pi_i(T, t)$ are representable by group schemes of finite type over $Spec(\mathbf{C})$, which are required to be affine for $i = 1$ and vector spaces (i.e. affine unipotent abelian) for $i \geq 2$.

(This is the “very presentability” condition of [41] under the additional hypothesis of connectedness; in the non-connected case the definition is more complicated and that is basically the subject of the paper [41].)

We now choose $\mathcal{P}$ for our shape theory to be the $n + 1$ -category of connected very presentable n -stacks of groupoids. Suppose $\mathcal{F}$ is an n -stack on $Sch/\mathbf{C}$. The *shape* of $\mathcal{F}$ is defined as the $n + 1$ -functor from the $n + 1$ -category of connected very presentable n -stacks T , to the $n + 1$ -category $nSTACK$ of all n -stacks, given by the formula

$$Shape(\mathcal{F})(T) := Hom(\mathcal{F}, T).$$

This contains all information about $\mathcal{F}$ which one can extract by looking at G -torsors over $\mathcal{F}$ and cohomology of associated vector bundles.

In many cases, $Hom(\mathcal{F}, T)$ will be a geometric or locally geometric n -stack [42]. For example in the case $\mathcal{F} = X_{Dol}$ we look at below, $Hom(X_{Dol}, T)$ will be locally geometric. In these cases the shape of $\mathcal{F}$ may be considered as an $n + 1$ -functor from connected very presentable n -stacks to the $n + 1$ -category of (locally) geometric n -stacks, sitting inside $nSTACK$.

Examples

We explain how to understand the structure of connected very presentable T . First is the simply connected case. Here T is given by a Postnikov tower where the stages are of the form $K(\mathcal{O}^a, m)$. The only question is how they are put together. The fibration

$$K(\mathcal{O}^a, m) \rightarrow \tau_{\leq m} T \rightarrow \tau_{\leq m-1} T$$

is classified by a map

$$\tau_{\leq m-1} T \rightarrow K(\mathcal{O}^a, m+1),$$

in other words $\tau_{\leq m} T$ is the pullback by this map of the standard fibration

$$K(\mathcal{O}^a, m) \rightarrow * \rightarrow K(\mathcal{O}^a, m+1).$$

We can write

$$\tau_{\leq m} T = \tau_{\leq m-1} T \times_{K(\mathcal{O}^a, m+1)} *.$$

The classifying map is a class in $H^{m+1}(\tau_{\leq m-1} T, \mathcal{O}^a)$. In turn, this cohomology can be “calculated” by the Leray-Serre spectral sequence applied to the previous part of the Postnikov tower for $\tau_{\leq m-1} T$. The basic pieces that we need to know are the cohomology of the Eilenberg-MacLane spaces. These are given by the *Breen calculations* [8] [9], which we recall in Appendix I (giving a relative version). For the present discussion the answer is that $H^*(K(\mathcal{O}^a, m), \mathcal{O})$ is a graded-symmetric algebra on $\mathcal{O}^a$ in degree m . Note that this answer is the same as the classical answer for rational cohomology of rational Eilenberg-MacLane spaces $H^*(K(\mathbf{Q}^a, m), \mathbf{Q})$.

As we shall explain below and also in Appendix II, if Y is a finite simply-connected CW -complex then we obtain a 1-connected very presentable $T = Y \otimes \mathbf{C}$ whose homotopy group sheaves are $\pi_i(Y, y) \otimes_{\mathbf{Q}} \mathcal{O}$.

We now look at connected but not simply connected very presentable T . Note that since T is connected, we can choose a basepoint $t \in T(\text{Spec}(\mathbf{C}))$. Let $G := \pi_1(T, t)$; by hypothesis it is an affine algebraic group scheme over $\mathbf{C}$. We have a fiber sequence

$$T' \rightarrow T \rightarrow K(G, 1)$$

where T' is simply connected very presentable. The homotopy group sheaves $\pi_i(T', t) = \pi_i(T, t)$ are vector spaces $\mathcal{O}^a$ for $i \geq 2$; but notice also that G acts on these vector spaces. The action is an action of sheaves on the site $Sch/\mathbf{C}$ so it is automatically algebraic; we can write $\pi_i(T', t) = V^i$ with V^i a linear representation of G .

The same Postnikov tower discussion as above, works here. The only difference is that in calculating the cohomology of the $\tau_{\leq m} T$ we may have coefficients which are linear representations of G , and at the end we get down to a step where we have to calculate $H^i(K(G, 1), V)$. If G is reductive, this “algebraic cohomology” vanishes for $i \geq 1$, whereas if G is unipotent then it is equal to the Lie algebra cohomology.

A simple example of non-simply connected T may be obtained as follows. Start with a linear algebraic group G with a linear representation V , corresponding to a local system $\underline{V} \rightarrow K(G, 1)$. Use the notational shorthand

$$K(V/G; n) := K(\underline{V}/K(G, 1), n).$$

We have a fibration sequence

$$K(V, n) \rightarrow K(V/G, n) \rightarrow K(G, 1).$$

Maps $\mathcal{F} \rightarrow T$ correspond to pairs (E, η) where E is a G -torsor over $\mathcal{F}$ and $\eta \in H^n(\mathcal{F}, E \times^G V)$.

An advantage of the nonabelian cohomological formulation of the secondary Kodaira-Spencer classes we define is that the definition works for cohomology with coefficients in a connected very presentable T , even non-simply connected. However, for the calculation we will do, we look at a particular simply-connected T (the “complexified 2-sphere” $S^2 \otimes \mathbf{C}$).

Representability

Under certain circumstances, basically when the shape of $\mathcal{F}$ is simply connected and has reasonable cohomology sheaves, then $\text{Shape}(\mathcal{F})$ is *representable*. By this we mean that there is a morphism $\mathcal{F} \rightarrow \Sigma$ from $\mathcal{F}$ to a very presentable n -stack Σ such that for any other very presentable n -stack T we have

$$\text{Hom}(\Sigma, T) \xrightarrow{\cong} \text{Hom}(\mathcal{F}, T).$$

For example we have the following precise statement.

Theorem 2.1 *Suppose $\mathcal{F}$ is an n -stack on $\text{Sch}/\mathbf{C}$ such that for any affine algebraic group G ,*

$$K(G, 1) \xrightarrow{\cong} \text{Hom}(\mathcal{F}, K(G, 1)).$$

Suppose that the $H^i(\mathcal{F}, \mathcal{O})$ are representable by finite dimensional vector spaces. Then there is a morphism $\mathcal{F} \rightarrow \Sigma$ to a 1-connected very presentable n -stack Σ , such that for any connected very presentable n -stack T we have

$$\text{Hom}(\Sigma, T) \xrightarrow{\cong} \text{Hom}(\mathcal{F}, T).$$

The proof will be given in Appendix II.

On the other hand, for n -stacks $\mathcal{F}$ whose shape is not 1-connected, the shape will not in general be representable. For example, suppose W is a finite CW complex (considered as a constant n -stack) such that $\pi_1(W) = \Gamma := \mathbf{Z}$ and such that some $\pi_i(W) \otimes \mathbf{Q}$ is a $\mathbf{Q}[\Gamma]$ -module which is not completely torsion (hence infinite-dimensional over $\mathbf{Q}$). Then $\text{Shape}(W)$ is not representable. Indeed, the infinite dimensionality of $\pi_i(W) \otimes \mathbf{Q}$ is seen by the shape, since all irreducible representations of Γ are finite (1-) dimensional.

3. Nonabelian Dolbeault cohomology

Suppose X is a smooth quasiprojective variety. Recall that one defines the *Dolbeault cohomology* of X as the hypercohomology of the trivial complex Ω_X with differential equal to 0:

$$H_{Dol}^i(X) := \mathbf{H}^i(X, \Omega_X^0 \xrightarrow{0} \dots) = \bigoplus_{p+q=i} H^q(X, \Omega_X^p)$$

(generally speaking this is only motivated by topology when X is projective; but we make the notation for quasiprojective X too, for use in Mayer-Vietoris arguments).

A nonabelian version for H^1 may be defined by setting $H_{Dol}^1(X, G)$ equal to the moduli stack of Higgs principal G -bundles ([22] [23] [36] [37]) (P, θ) . One can impose semistability conditions and vanishing of Chern classes to get a version more closely related to topology, but we don't need that for the present algebraic discussion.

If (P, θ) is a principal Higgs bundle and if V is a representation of G then we obtain an associated Higgs bundle (E, θ) . Recall that we define the *Dolbeault cohomology of (E, θ)* as the hypercohomology of the *Dolbeault complex* (cf [36])

$$\dots \xrightarrow{\theta} E \otimes_{\mathcal{O}_X} \Omega_X^i \xrightarrow{\theta} \dots,$$

$$H_{Dol}^i(X, (E, \theta)) := \mathbf{H}^i(X, (E \otimes_{\mathcal{O}_X} \Omega_X, \theta)).$$

We present a way of unifying these definitions into a notion of *nonabelian Dolbeault cohomology*. For nonabelian H^1 the present interpretation was explained in [40].

Let $\widehat{TX}$ denote the formal completion of TX along the zero section. Considered as a presheaf on $Sch/\mathbf{C}$ it associates to any $\mathbf{C}$ -scheme Y , the set of maps $Y \rightarrow TX$ which map the underlying reduced subscheme Y^{red} to the zero-section $X \subset TX$.

Define the 1-stack X_{Dol} to be the relative $K(\widehat{TX}/X, 1)$, i.e. the relative classifying stack for the group scheme $\widehat{TX} \rightarrow X$.

A variant which is technically easier to work with is

$$X_{UDol} := K(TX/X, 1).$$

The U in the notation stands for “unipotent”: as we shall see below (Proposition 3.1), a morphism $X_{UDol} \rightarrow K(G, 1)$ corresponds to a principal Higgs bundle with structure group G , over X , such that the Higgs field is a section of unipotent elements of the Lie algebra of G . However, for morphisms to simply connected T , we can safely replace X_{Dol} by X_{UDol} and for the purposes of the present paper this is what we shall do.

If T is an n -stack then we define the *nonabelian Dolbeault cohomology of X with coefficients in T* to be the n -stack (of n -groupoids)

$$Hom(X_{Dol}, T).$$

Of course this includes the classical abelian case, when we take $T = K(\mathcal{O}, n)$ where $\mathcal{O}$ denotes the structural sheaf, equal to $\mathbf{G}_a$, represented by the affine line. It also includes the somewhat classical case of nonabelian Dolbeault H^1 with coefficients in a group scheme G .

More generally we will be most interested in the case where T is a connected very presentable n -stack. The calculation of the present paper involves a 1-connected T , so it doesn't refer to the case of Higgs principal G -bundles or Dolbeault cohomology with coefficients in Higgs bundles. These aspects are only presented to show the unified character of the definition.

Proposition 3.1 *We have that*

$$\pi_0(\mathrm{Hom}(X_{Dol}, K(\mathcal{O}, n))) = H_{Dol}^n(X, \mathbf{C})$$

is the usual Dolbeault cohomology of X . The same holds for X_{UDol} . If G is an affine algebraic group scheme then

$$\mathrm{Hom}(X_{Dol}, K(G, 1)) = \mathcal{M}_{Dol}(X, G)$$

is the moduli 1-stack of principal Higgs bundles with structure group G . On the other hand,

$$\mathrm{Hom}(X_{UDol}, K(G, 1))$$

is the moduli 1-stack of principal Higgs bundles (P, θ) with structure group G , such that for every $x \in X$ the element $\theta_x \in \mathrm{ad}(P)_x \cong \mathfrak{g}$ is a unipotent element of the Lie algebra $\mathfrak{g}$ of G (this condition is of course independant of the isomorphism $\mathrm{ad}(P)_x \cong \mathfrak{g}$ chosen).

Proof: Let X_{Dol}^{fc} denote the formal category defined in [40] which gives the 1-stack X_{Dol} . We have

$$\mathrm{Ob} X_{Dol}^{\mathrm{fc}} = X,$$

and

$$\mathrm{Mor}(X_{Dol}^{\mathrm{fc}}) = \widehat{TX} \rightarrow X \hookrightarrow X \times X$$

(the morphism object lies over the diagonal in $X \times X$). The composition of morphisms is just addition in $\widehat{TX}$ (which is a formal group scheme over X). This formal groupoid gives in an obvious way a presheaf of groupoids on $Sch/\mathbf{C}$, whose associated stack is X_{Dol} . A morphism $X_{Dol} \rightarrow K(G, 1)$ is the same thing as a G -torsor over X_{Dol}^{fc} , which in turn is the same thing as a principal G -bundle P over X together with action of the formal group scheme $\widehat{TX}$. The action may be interpreted as a morphism of sheaves of groups over X ,

$$\widehat{TX} \rightarrow \mathrm{Ad}(P).$$

Since the domain is formal, this is the same thing as a morphism of Lie algebras over X ,

$$\theta : TX \rightarrow ad(P).$$

This proves the second statement. For the third statement, the same proof works but with $\widehat{TX}$ replaced by TX . Note that TX is a unipotent group scheme over X , so

$$TX \rightarrow Ad(P)$$

corresponds to a morphism of Lie algebras

$$\theta : TX \rightarrow ad(P)$$

with image in the unipotent elements of $ad(P)$.

For the first statement, one way to proceed is to notice that the cohomology of X_{Dol} with coefficients in $\mathcal{O}$ is the same as the cohomology of the formal category X_{Dol}^{fc} as considered by Berthelot [5] and Illusie [24]. From those references, one gets a generalized de Rham complex calculating the cohomology, which in our case is seen to be exactly the Dolbeault complex.

We also need to prove that the morphism $X_{Dol} \rightarrow X_{UDol}$ induces an isomorphism on cohomology. For this it suffices to look locally over X , so we can assume that TX is trivial. Thus it suffices to prove that if V is a vector space of dimension n then the morphism

$$K(\widehat{V}, 1) \rightarrow K(V, 1)$$

induces an isomorphism of cohomology. This morphism has homotopy fiber the sheaf V_{DR} defined by $V_{DR}(Y) = V(Y^{\text{red}})$ (cf [39]). The cohomology of V_{DR} with coefficients in $\mathcal{O}$ is the algebraic de Rham cohomology of V ([39] Theorem 6.2) which is trivial because V is an affine space.

Since the statement of the first part for X_{UDol} is what we actually use, we indicate a somewhat more elementary proof. Let V be a vector space of dimension $n = \dim(X)$. We have a natural (split) extension

$$1 \rightarrow V \rightarrow G \rightarrow GL(V) \rightarrow 1$$

which gives a fiber sequence of 1-stacks

$$K(V, 1) \rightarrow K(G, 1) \xrightarrow{p} K(GL(V), 1).$$

Let $C^\bullet = R^*p_*(\mathcal{O})$. Using the Breen calculations [9] [39], which we recall in Theorem 9.2 in Appendix I below, it is easy to see that

$$H^i(C^\bullet) = \bigwedge^i (V^*).$$

However, the fact that $GL(V)$ is a reductive group implies that all of the algebraic group cohomology, in other words the cohomology of $K(GL(V), 1)$ with coefficients in any local system associated to a representation of $GL(V)$, vanishes except in degree 0. Therefore the invariants of the complex $C^\bullet$ vanish so

$$C^\bullet \sim \bigoplus_i \bigwedge^i (V^*).$$

Now notice that the vector bundle $TX \rightarrow X$ corresponds to a map $X \rightarrow K(GL(V), 1)$ and

$$X_{UDol} = K(G, 1) \times_{K(GL(V), 1)} X.$$

Let $q : X_{UDol} \rightarrow X$ denote the projection; then $Rq_*\mathcal{O}$ is the pullback of $C^\bullet$, hence it splits as

$$Rq_*\mathcal{O} = \bigoplus_i \Omega_X^i.$$

The cohomology of X_{UDol} with coefficients in $\mathcal{O}$ is equal to the hypercohomology of X with coefficients in this complex, which is the Dolbeault cohomology. $\square$

Remark: We have the following which are sometimes useful:

$$\pi_i(\text{Hom}(X_{Dol}, K(\mathcal{O}, n))) = H_{Dol}^{n-i}(X, \mathbf{C}).$$

Proposition 3.2 *Suppose (P, θ) is a Higgs principal G -bundle on X corresponding to a map $X_{Dol} \rightarrow K(G, 1)$. Suppose V is a representation of G and let (E, θ) be the associated Higgs bundle. Then the cohomology of X_{Dol} with coefficients in the local system $(P, \theta)^*(V)$ is naturally isomorphic to the Dolbeault cohomology $H_{Dol}^n(X, (E, \theta))$. The same works for a family of principal bundles parametrized by a base scheme S .*

Proof: As before, we interpret (P, θ) as a G -torsor over the formal category X_{Dol}^{fc} which defines the stack X_{Dol} . Associated to the representation V we get a local system over the formal category, and its cohomology (which is that mentioned in the statement of the proposition) is calculated by a de Rham complex (cf [5] [24]). This de Rham complex is exactly the Dolbeault complex for (E, θ) . $\square$

Corollary 3.3 *Suppose T is a connected very presentable n -stack such that $\pi_1(T)$ is a unipotent group scheme over $\text{Spec}(\mathbf{C})$. Then the morphism*

$$\text{Hom}(X_{UDol}, T) \rightarrow \text{Hom}(X_{Dol}, T)$$

is an equivalence.

Proof: This follows immediately from the previous propositions using a Postnikov tower for T , and noting that if G is a unipotent group scheme then all principal G -Higgs bundles satisfy the condition that the Higgs field be a section of unipotent elements. $\square$

Cohomology classes of subschemes

A particular detail of (usual) Dolbeault cohomology which enters into our calculations is the cohomology class of a subscheme. Suppose X is a smooth projective variety and suppose that $Z \subset X$ is a subscheme of codimension d . Then we obtain the class

$$[Z] \in H^d(X, \Omega_X^d) \subset H_{Dol}^{2d}(X).$$

In fact this class has a canonical trivialization over $U := X - Z$, in other words we are actually given a lifting to

$$[Z] \in H^d((X, U), \Omega_X^d) \subset H^{2d}((X_{Dol}, U_{Dol}), \mathcal{O}).$$

Suppose for example $d = 1$. Take an open covering $X = \bigcup U_i$ with $U_0 = U = X - Z$, and the remaining U_i affine with defining equations $z_i \in \mathcal{O}(U_i)$ having zeros of order one along Z and nonvanishing elsewhere. Put

$$g_{ij} := \frac{dz_i}{z_i} - \frac{dz_j}{z_j}.$$

This gives a 1-cocycle which determines the cohomology class $[Z]$. Since $U_0 = X - Z$ is a part of the covering, it gives a class in the cohomology of the pair $(X, X - Z)$.

The case of higher codimension is treated similarly. The only thing we need to know is that if $\lambda, \lambda' \in \mathcal{O}(X)$ are regular functions such that $\lambda|_Z = \lambda'|_Z$, then $\lambda[Z] = \lambda'[Z]$. In particular if P is a point then $\lambda[P] = \lambda(P)[P]$.

4. Group actions

Suppose W is a sheaf of groups on $Sch/\mathbf{C}$, and suppose R is an n -stack on $Sch/\mathbf{C}$. Then an *action of W on R* is a morphism

$$\rho : A \rightarrow K(W, 1)$$

together with an identification of the fiber $\rho^{-1}(0)$ (by which we mean the homotopy fiber product $\{0\} \times_{K(W, 1)} A$) with R ,

$$R \cong \rho^{-1}(0).$$

To put this more briefly, an action of W on R is a fiber sequence

$$R \rightarrow A \rightarrow K(W, 1).$$

If T is another n -stack, and if ρ is an action of W on R then we obtain an associated action $\text{Hom}(\rho, T)$ of W on $\text{Hom}(R, T)$. It is given by the relative Hom ,

$$\text{Hom}(A/K(W, 1), T) \rightarrow K(W, 1),$$

whose fiber is naturally identified with $\text{Hom}(R, T)$.

Example 1: If W acts on a sheaf of sets R in the usual sense, then this can be interpreted as an action in the above sense. The stack A (which in this case is a 1-stack) is the stack-theoretical quotient R/W with its canonical principal W -bundle $R \rightarrow R/W$ which corresponds to a morphism $R/W \rightarrow K(W, 1)$.

Example 2: The group $\mathbf{G}_m$ acts on $\widehat{TX}$ (resp. TX) over X by scalar multiplication. Therefore $\mathbf{G}_m$ acts on X_{Dol} (resp. X_{UDol}) and on $\text{Hom}(X_{Dol}, T)$ (resp. $\text{Hom}(X_{UDol}, T)$). In the case $T = K(G, 1)$ this is the usual action of $\mathbf{G}_m$ on the moduli stack of principal Higgs bundles [23] [36] [37]. In the case $T = K(\mathcal{O}, n)$ this action gives the decomposition of $H_{Dol}^i(X)$ into pieces $H^q(X, \Omega_X^p)$.

Example 3: A different example, more closely related to what we are interested in, is the following. Suppose

$$1 \rightarrow V \rightarrow E \rightarrow W \rightarrow 1$$

is an exact sequence of sheaves of groups on $Sch/\mathbf{C}$. This gives a fibration sequence

$$K(V, 1) \rightarrow K(E, 1) \rightarrow K(W, 1),$$

hence by definition it is an action of W on $K(V, 1)$.

More generally if X is a sheaf of sets and if $V \rightarrow X$ is a sheaf of groups over X , then we can look at the relative Eilenberg-MacLane stack $K(V/X, 1) \rightarrow X$. Suppose

$$1 \rightarrow V \rightarrow E \rightarrow p^*(W) \rightarrow 1$$

is an exact sequence of sheaves of groups on X (where $p : X \rightarrow *$ denotes the projection). Then we obtain a fibration sequence

$$K(V/X, 1) \rightarrow K(E/X, 1) \rightarrow K(W, 1),$$

the latter map being the composition of the map induced by the second map in the exact sequence, with the projection

$$K(p^*W/X, 1) = K(W \times X/X, 1) = K(W, 1) \times X \rightarrow K(W, 1).$$

Thus, again by definition, our exact sequence corresponds to an action of W on $K(V/X, 1)$ (lying over the trivial action of W on X).

Secondary classes

We first discuss classifying spaces. If R is an n -stack then $Aut(R)$ is an n -stack with “group” structure, more precisely with the structure of a loop space. To make this statement precise, we construct a pointed $n + 1$ -stack $B Aut(R)$ with basepoint denoted 0 , with an equivalence

$$\Omega(B Aut(R), 0) \cong Aut(R).$$

Construct the presheaf of spaces $B^{\text{pre}} Aut(R)$ as the realization of a simplicial n -stack $B.Aut(R)$ defined as follows: let $\bar{I}^{(i)}$ be the 1-category with $i+1$ isomorphic objects $0, \dots, i$ (which we shall call “vertices”); then set $B_i Aut(R)$ equal to the n -stack of morphisms $\bar{I}^{(i)} \rightarrow n\text{STACK}$ sending the vertices to the object $R \in n\text{STACK}$. Note that the component $B_i Aut(R)$ is homotopic to $Aut(R) \times \dots \times Aut(R)$. This simplicial n -stack may be interpreted as a presheaf of $n + 1$ -categories. Note that $Aut(R)$ is defined to be the n -stack of morphisms $\bar{I} \rightarrow n\text{STACK}$ sending $0, 1$ to R , and it is a stack of n -groupoids. It may safely be confused with a presheaf of n -truncated spaces, and $B.Aut(R)$ becomes a presheaf of simplicial spaces. The component simplicial spaces satisfy Segal’s condition (cf [35] [46] [43] [45]) so if we set $B^{\text{pre}} Aut(R)$ equal to the realization into a presheaf of spaces then Segal’s Proposition 1.5 [35] implies that the natural map

$$Aut(R) = B_1 Aut(R) \rightarrow \Omega(B^{\text{pre}} Aut(R), 0)$$

is an equivalence (object-by-object). Finally, let $B Aut(R)$ be the $n + 1$ -stack associated to $B^{\text{pre}} Aut(R)$.

If R is an n -stack of groupoids, the locally constant fibrations with fiber R

$$R \rightarrow E \rightarrow S$$

are classified exactly by maps $S \rightarrow B Aut(R)$. In other words, given S and R , the $n + 1$ -category of such fiber sequences is equivalent to the $n + 1$ -category $Hom(S, B Aut(R))$.

In particular, an action of a sheaf of groups W on an n -stack of groupoids R is the same thing as a morphism

$$f : K(W, 1) \rightarrow B Aut(R).$$

This leads to the notion of *characteristic classes* for the action: if $\mathcal{G}$ is a sheaf of groups and if $c \in H^i(B Aut(R), \mathcal{G})$ then for any action we can pull back c to obtain a class in $H^i(K(W, 1), \mathcal{G})$.

One can also obtain *secondary invariants*, which only become defined when some primary invariants vanish. For example, suppose $c \in H^i(B Aut(R), \mathcal{G})$ is a cohomology class corresponding to a map

$$c : B Aut(R) \rightarrow K(\mathcal{G}, i).$$

Suppose that $f : K(W, 1) \rightarrow B \operatorname{Aut}(R)$ is an action of W on R , such that $c \circ f : K(W, 1) \rightarrow K(\mathcal{G}, i)$ is trivial, i.e. homotopic to the constant map at the basepoint. Let $\operatorname{Fib}(c)$ denote the homotopy fiber of the morphism c over 0. A choice of trivialization ψ of $c \circ f$ gives rise to a morphism

$$f_\psi : K(W, 1) \rightarrow \operatorname{Fib}(c).$$

The choices of trivialization ψ are, up to homotopy, classified by $H^{i-1}(K(W, 1), \mathcal{G})$. If c' is a class in $H^j(\operatorname{Fib}(c), \mathcal{G}')$ then the composition $c' \circ f_\psi$ gives a class in $H^j(K(W, 1), \mathcal{G}')$. This is (a typical example of) a secondary characteristic class for the action.

We don't get any further into the general theory of secondary classes, because insofar as the formalism is concerned, there is no difference between the present case of n -stacks and the classical topological version of the theory.

Instead, we give a more explicit description of the secondary classes that we are interested in. These concern only classes of degrees 1 and 2 on $B \operatorname{Aut}(R)$, so a more concrete discussion is possible. The first step is to replace R by its truncation down to a 1-stack, denoted $\tau_{\leq 1}R$. We fix a point $r \in R(\operatorname{Spec}(\mathbf{C}))$ (and denote also by r its image in $\tau_{\leq 1}(R)$). Note that the “group” $\operatorname{Aut}(R)$ acts on $\tau_{\leq 1}R$. The first of the primary classes comes from a morphism

$$\operatorname{Aut}(R) \rightarrow \operatorname{Aut}(\pi_0 R).$$

On the left is an actual sheaf of groups $\mathcal{G}$, and this map corresponds to a class in $H^1(B \operatorname{Aut}(R), \mathcal{G})$. Suppose that we have an action by W such that this class vanishes. This implies that W fixes the point $r \in \pi_0(R)(\operatorname{Spec}(\mathbf{C}))$. (Vanishing of this class actually implies that W fixes every point but for what follows we only need that it fixes r). Let R^r denote the component of R containing r , more precisely it is the fiber product

$$R^r := * \times_{\pi_0(R)} R.$$

Under our preliminary vanishing hypothesis, we obtain an action of W on R^r .

Now $\tau_{\leq 1}(R^r) = K(G, 1)$ where $G = \pi_1(R, r)$ (the point r is a basepoint defined over $\operatorname{Spec}(\mathbf{C})$ giving this trivialization of the 1-gerb $\tau_{\leq 1}(R^r)$ —the basepoint r corresponds to $0 \in K(G, 1)$).

For any group G we have a fibration sequence

$$K(\operatorname{Center}(G), 2) \rightarrow B \operatorname{Aut}(K(G, 1)) \rightarrow K(\operatorname{Out}(G), 1).$$

In our case above, W acts on R^r , hence on $\tau_{\leq 1}R^r = K(G, 1)$. The map

$$K(W, 1) \rightarrow B \operatorname{Aut}(\tau_{\leq 1}R^r) = B \operatorname{Aut}(K(G, 1))$$

thus projects first of all to a morphism of groups

$$W \rightarrow \operatorname{Out}(\pi_1(R, r)).$$

This is in a certain sense again a primary invariant. Suppose that this invariant vanishes. Then, given that

$$K(W, 1) \rightarrow K(Out(G), 1)$$

is a pointed map, there is a canonical homotopy of this map to the constant map at the basepoint, so we canonically can identify our map

$$K(W, 1) \rightarrow B Aut(\tau_{\leq 1} R^r)$$

as a map

$$K(W, 1) \rightarrow K(Center \pi_1(R, r), 2),$$

in other words as a class in $H^2(K(W, 1), Center)$ where *Center* is the center of $\pi_1(R, r)$. This is the secondary class we will be interested in calculating below.

We can describe the secondary class a bit more concretely in the following way. Suppose α, β are elements of W thought of as paths in $K(W, 1)$. Choose trivializations of the fibration

$$R \rightarrow A \rightarrow K(W, 1)$$

above the paths α and β . These trivializations lead in particular to liftings of our paths starting with the basepoint r , and ending at points we denote $\alpha^*(r)$ and $\beta^*(r)$. By the hypothesis of vanishing of primary classes, $\alpha^*(r)$ and $\beta^*(r)$ are homotopic to r . Thus we can choose homotopies which we denote h_α and h_β (these are paths in R joining $\alpha^*(r)$ resp. $\beta^*(r)$ to r). Applying the trivializations chosen above to these paths we obtain paths in R

$$\alpha^*(h_\beta) : \alpha^* \beta^*(r) \rightarrow \alpha^*(r),$$

$$\beta^*(h_\alpha) : \beta^* \alpha^*(r) \rightarrow \beta^*(r).$$

Finally, the commutativity of W means that there is a torus obtained by attaching a 2-cell along the commutator of α and β . Lifting this 2-cell (more precisely, trivializing the family above this 2-cell) provides a path between $\alpha^* \beta^*(r)$ and $\beta^* \alpha^*(r)$. Combining these all together we get a 5-sided loop based at r ; this is the secondary class evaluated on $\alpha \wedge \beta$. The loop is in the center of $\pi_1(R, r)$ because of the fact that we chose trivializations of the family over our paths (or 2-cell) rather than just liftings starting at the basepoint.

A slightly different and more geometric way of looking at this is to look at the torus in $K(W, 1)$ given by the commutator 2-cell for α and β . Attaching two 2-cells to the torus, one along α and the other along β , gives a 2-sphere. The vanishing of the primary classes means that the fibration $A \rightarrow K(W, 1)$ can be extended across these new 2-cells, so we get a family over S^2 . Trivializing over the northern and southern hemispheres, the family is determined by a morphism from the equator to $Aut(R)$: this element of $\pi_1(Aut(R)) = Center$ is the image of $\alpha \wedge \beta$ under the secondary class.

5. The secondary Kodaira-Spencer map

We now come to the situation which gives a “secondary Kodaira-Spencer map”. Suppose $f : X_S \rightarrow S$ is a smooth projective morphism. Fix a basepoint $s \in S$ and denote by X the fiber of X_S over s . Define

$$W := T(S)_s$$

which is a vector space, thus unipotent abelian group scheme, considered as a sheaf of groups on $Sch/\mathbf{C}$.

We have an exact sequence (of unipotent abelian group schemes i.e. vector bundles over X)

$$0 \rightarrow TX \rightarrow T(X_S)|_X \rightarrow f^*(T(S)_s) \rightarrow 0.$$

This is an action of $W = T(S)_s$ on the stack $K(TX/X, 1) = X_{UDol}$. In particular, for any n -stack T we obtain an action of W on $Hom(X_{UDol}, T)$.

The primary invariant in this situation is an action of W on the sheaf of sets $\pi_0 Hom(X_{UDol}, T)$. For example, if $T = K(\mathcal{O}, n)$ then by Proposition 3.1

$$\pi_0 Hom(X_{UDol}, T) = H_{Dol}^n(X) = \bigoplus_{p+q=n} H^q(X, \Omega_X^p)$$

and we obtain an action of $W = T(S)_s$ on $H_{Dol}^n(X)$. This action is of course just the usual Kodaira-Spencer map which decomposes into components

$$T(S)_s \rightarrow Hom\left(H^q(X, \Omega_X^p), H^{q+1}(X, \Omega_X^{p-1})\right).$$

To obtain secondary invariants, we proceed as described above, using vanishing of the primary invariants if we want to (but bearing in mind that the secondary invariants will then only be defined when the primary invariants vanish). For example, suppose that T is an n -stack and suppose $\eta \in Hom(X_{UDol}, T)$ such that the point $[\eta] \in \pi_0 Hom(X_{UDol}, T)$ is fixed by the action of W . Then W acts on the connected n -stack $Hom^\eta(X_{UDol}, T)$ which is the connected component containing η . We obtain a morphism

$$K(W, 1) \rightarrow B Aut(Hom^\eta(X_{UDol}, T)),$$

and as remarked above, cohomology classes on the right can be pulled back to give classes on $K(W, 1)$. At this point we refer to the Breen calculations [9]. In Appendix II we prove a relative version in characteristic zero which was stated in [39]; the reader may refer there for the general statement. In our case, as W is represented by a finite-dimensional vector space (in particular, $W \cong \mathcal{O}^a$) we have

$$H^i(K(W, 1), \mathcal{O}) = \bigwedge^i W^*$$

where $W^* = \text{Hom}(W, \mathcal{O})$ is the sheaf represented by the dual vector space.

To get down to the concrete example we would like to consider, we boil things down a bit farther, following the discussion at the end of the previous section. Namely instead of looking at the full $\text{Hom}^\eta(X_{UDol}, T)$ we truncate it down to a 1-stack (which is connected, too) by looking at

$$\tau_{\leq 1} \text{Hom}^\eta(X_{UDol}, T).$$

In the very presentable case, this automatically has a base point over $\text{Spec}(\mathbf{C})$ (but also we have chosen a basepoint η) so it is equivalent to something of the form $K(G, 1)$ with

$$G = \pi_1(\text{Hom}(X_{UDol}, T), \eta).$$

Again, in the very presentable case (i.e. if T is very presentable which implies the same for $\text{Hom}^\eta(X_{UDol}, T)$) then G is an affine algebraic group. In our example below G will itself be a vector space. The first invariant is

$$W \rightarrow \text{Out}(\pi_1(\text{Hom}(X_{UDol}, T), \eta)).$$

Suppose that this invariant vanishes. Then we get a map

$$K(W, 1) \rightarrow K(\text{Center } \pi_1(\text{Hom}(X_{UDol}, T), \eta), 2),$$

which may be interpreted as a class in $H^2(K(W, 1), \text{Center})$ where Center is the center of $\pi_1(\text{Hom}(X_{UDol}, T), \eta)$. If T is very presentable then this center will be an affine abelian group scheme. This class in H^2 is the class we are interested in calculating below.

The secondary class can be described concretely by choosing homotopies trivializing the primary classes and combining them together using the commutativity homotopy for the action of W , as at the end of the previous section.

6. The complexified 2-sphere

We discuss in more detail the example of a 3-stack T for which we make our calculation. Recall from “standard topology” how to describe the rational homotopy type of S^2 . The only nontrivial stages in the Postnikov tower are $K(\mathbf{Q}, 2)$ and $K(\mathbf{Q}, 3)$. Thus the rational homotopy type is described by the fibration sequence

$$K(\mathbf{Q}, 3) \rightarrow S^2 \otimes \mathbf{Q} \rightarrow K(\mathbf{Q}, 2).$$

In turn, the classifying space for fibrations with fiber $K(\mathbf{Q}, 3)$ is the base for the universal fibration

$$K(\mathbf{Q}, 3) \rightarrow * \rightarrow K(\mathbf{Q}, 4).$$

Thus the above fibration is determined by a morphism

$$K(\mathbf{Q}, 2) \rightarrow K(\mathbf{Q}, 4),$$

in other words a class $\sigma \in H^4(K(\mathbf{Q}, 2), \mathbf{Q})$. The classical calculations give

$$H^{2m}(K(\mathbf{Q}, 2), \mathbf{Q}) = \text{Sym}^m(\mathbf{Q}).$$

In particular

$$H^4(K(\mathbf{Q}, 2), \mathbf{Q}) \cong \text{Sym}^2(\mathbf{Q}) = \mathbf{Q}.$$

Up to change of basis element for π_3 , there are only two possibilities: either $\sigma = 0$ or $\sigma \neq 0$. The case $\sigma = 0$ corresponds to the direct product $K(\mathbf{Q}, 2) \times K(\mathbf{Q}, 3)$ but in this case H^3 would be nonzero, whereas $H^3(S^2 \otimes \mathbf{Q}, \mathbf{Q}) = 0$. Therefore we must have $\sigma \neq 0$, in other words σ is the cup product $\eta \cup \eta$ of the canonical class $\eta \in H^2(K(\mathbf{Q}, 2), \mathbf{Q})$ with itself.

In view of this discussion, we can do the same with very presentable stacks. We will construct $T = S^2 \otimes \mathbf{C}$ with two stages $K(\mathcal{O}, 2)$ and $K(\mathcal{O}, 3)$ in the Postnikov tower. We construct it as the pullback of the universal fibration

$$K(\mathcal{O}, 3) \rightarrow * \rightarrow K(\mathcal{O}, 4)$$

by a morphism

$$\sigma : K(\mathcal{O}, 2) \rightarrow K(\mathcal{O}, 4).$$

The Breen calculations say that the cohomology of $K(\mathcal{O}, n)$ with coefficients in $\mathcal{O}$ has the same answer as the cohomology of $K(\mathbf{Q}, n)$ with coefficients in $\mathbf{Q}$. In other words,

$$H^{2m}(K(\mathcal{O}, 2), \mathcal{O}) = \text{Sym}_{\mathcal{O}}^m(\mathcal{O}) \cong \mathcal{O}$$

and

$$H^4(K(\mathcal{O}, 2), \mathcal{O}) = \mathcal{O}.$$

Let σ be the generator of $\mathcal{O}(\text{Spec } \mathbf{C}) = \mathbf{C}$. This gives a map

$$\sigma : K(\mathcal{O}, 2) \rightarrow K(\mathcal{O}, 4).$$

In terms of cohomology operations, a morphism $\mathcal{F} \rightarrow K(\mathcal{O}, 2)$ corresponds to a cohomology class $\eta \in H^2(\mathcal{F}, \mathcal{O})$ and the composition of such a map with σ corresponds to the cup-product square $\eta \cup \eta \in H^4(\mathcal{F}, \mathcal{O})$.

Now set

$$T := S^2 \otimes \mathbf{C} := K(\mathcal{O}, 2) \times_{K(\mathcal{O}, 4)} *,$$

(this choice of T shall be in vigor for the rest of the paper unless explicitly mentionned otherwise). For any n -stack $\mathcal{F}$, a morphism

$$\mathcal{F} \rightarrow T$$

corresponds to a pair (η, φ) where $\eta : \mathcal{F} \rightarrow K(\mathcal{O}, 2)$ and φ is a homotopy between

$$\sigma \circ \eta = \eta \cup \eta : \mathcal{F} \rightarrow K(\mathcal{O}, 4)$$

and the constant map at the basepoint

$$\underline{\sigma} : \mathcal{F} \rightarrow K(\mathcal{O}, 4).$$

Alternatively, φ may be thought of as a section of the fibration

$$K(\mathcal{O}, 3) \rightarrow \mathcal{F} \times_{K(\mathcal{O}, 4)} * \rightarrow \mathcal{F}.$$

A first remark is that for a given class $\eta \in H^2(\mathcal{F}, \mathcal{O})$ there exists a lifting to a map $\mathcal{F} \rightarrow T$ if and only if $\eta \cup \eta = 0$ in $H^4(\mathcal{F}, \mathcal{O})$. If this cup-product is zero so that there exists one lifting, then the fiber of the map

$$Hom(\mathcal{F}, T) \rightarrow Hom(\mathcal{F}, K(\mathcal{O}, 2))$$

over the point η is equivalent to $Hom(\mathcal{F}, K(\mathcal{O}, 3))$. This is because if there exists one lifting then we can choose a lifting to trivialize the fibration

$$\mathcal{F} \times_{K(\mathcal{O}, 4)} * \rightarrow \mathcal{F}$$

i.e. to make this fibration equivalent to

$$\mathcal{F} \times K(\mathcal{O}, 3) \rightarrow \mathcal{F}$$

and then the stack of other liftings is just $Hom(\mathcal{F}, K(\mathcal{O}, 3))$.

Example: We give an example of a topological space Y which admits no nonconstant maps to the actual 2-sphere S^2 but which admits maps to the complexified 2-sphere $S^2 \otimes \mathbf{C}$. Construct Y by taking a wedge of two 2-spheres and adding on a 4-cell via an attaching map

$$f : S^3 \rightarrow S^2 \vee S^2.$$

Note that

$$\pi_3(S^2 \vee S^2) = Sym^2 \pi_2(S^2 \vee S^2) = \mathbf{Z}^3,$$

and the class of the attaching map f determines the cup product. We can think of the class of f as a symmetric 2×2 matrix, which can be chosen arbitrarily. Choose the matrix

to be diagonal with (r, s) on the diagonal. The two obvious classes $e, f \in H^2(S^2 \vee S^2, \mathbf{Q})$ persist as classes in $H^2(Y, \mathbf{Q})$. Note that $H^4(Y, \mathbf{Q}) \cong \mathbf{Q}$ and we have the formulae

$$e \cup e = r, \quad f \cup f = s, \quad e \cup f = 0.$$

Now a map $Y \rightarrow S^2$ (or even to $S^2 \otimes \mathbf{Q}$) corresponds to a class

$$\eta = ae + bf \in H^2(Y, \mathbf{Q})$$

with $a, b \in \mathbf{Q}$, such that $\eta \cup \eta = 0$. The lifting to a map into S^2 is unique because in our case $H^3(Y, \mathbf{Q}) = 0$. However, if r and s are chosen to be relatively prime and having no square prime factors, then the equation

$$\eta \cup \eta = a^2 r + b^2 s = 0$$

doesn't have any nonzero solutions with $a, b \in \mathbf{Q}$ (the same holds if r and s have the same sign). Thus there are no nontrivial maps $Y \rightarrow S^2$. On the other hand, the above equation always has nonzero complex solutions, so there is always a nontrivial map $Y \rightarrow S^2 \otimes \mathbf{C}$.

7. Our example

We now come down to the example which we would like to calculate. Fix $T = S^2 \otimes \mathbf{C}$ as defined above. Let Z be a smooth projective surface with $H^1(Z, \mathcal{O}) = 0$ (hence $H^1(Z, \mathbf{C}) = H^3(Z, \mathbf{C}) = 0$). Let $P \in Z$ and let X be the blow-up of Z at P . Let $W \subset H^1(X, TX)$ be the rank two subspace of deformations of X corresponding to moving the point P which is blown up (thus canonically $W \cong T(Z)_P$). Then W acts on X_{UDol} , via the exact sequence

$$0 \rightarrow TX \rightarrow T(Tot)|_X \rightarrow W \otimes_{\mathbf{C}} \mathcal{O}_X \rightarrow 0$$

where Tot refers to the total space of the family. Let

$$R := Hom(X_{UDol}, T).$$

Then W acts on R and we will look at secondary classes for this action.

Because of our hypothesis $H_{Dol}^3(X) = 0$, the map $T \rightarrow K(\mathcal{O}, 2)$ induces an injection

$$\pi_0(R) \hookrightarrow H_{Dol}^2(X)$$

with the image the sheaf represented by the subscheme defined by the equation $\eta \cup \eta = 0$. The variation of Hodge structure of $H^2(X, \mathbf{C})$ parametrized by $P \in Z$ as we move the point which is blown up, is trivial. Thus the Kodaira-Spencer class is trivial, in other

words W acts trivially on $H_{Dol}^2(X)$ and hence it acts trivially on $\pi_0(R)$. If we fix $\rho = (\eta, \varphi) \in R(\text{Spec } \mathbf{C})$ then we obtain a secondary class

$$\kappa \in H^2(K(W, 1), \pi_1(R, \rho)).$$

The first task is to calculate $\pi_1(R, \rho)$ (which will be a group scheme over $\text{Spec}(\mathbf{C})$ since ρ is defined over $\text{Spec}(\mathbf{C})$). Recall that we have the fibration

$$R \rightarrow \text{Hom}(X_{UDol}, K(\mathcal{O}, 2))$$

whose fiber over a point in the target is either empty or else equivalent to $\text{Hom}(X_{UDol}, K(\mathcal{O}, 3))$. Thus the long exact sequence for this fibration gives

$$\mathbf{C} = H_{Dol}^0(X) \rightarrow H_{Dol}^2(X) \rightarrow \pi_1(R, \rho) \rightarrow H_{Dol}^1(X) = 0.$$

the first term being $\pi_2 \text{Hom}(X_{UDol}, K(\mathcal{O}, 2))$, the second term being $\pi_1 \text{Hom}(X_{UDol}, K(\mathcal{O}, 3))$, and so on.

Claim: The connecting morphism $H_{Dol}^0(X) \rightarrow H_{Dol}^2(X)$ in the above exact sequence is multiplication by a nonzero multiple of η .

Proof: For any n -stack Y we can define $\text{Hom}(Y, T)$ and look at the long exact sequence for the fibration

$$\text{Hom}(Y, T) \rightarrow \text{Hom}(Y, K(\mathcal{O}, 2)).$$

The connecting morphism will be functorial in Y . Apply this to $Y = T$ itself; then for any other Y (such as X_{UDol} considered in the claim) the connecting morphism for the long exact sequence at a basepoint $Y \rightarrow T$ is obtained by pulling back the connecting morphism for $\text{Hom}(T, T)$ at the identity map. In other words, looking at $Y = T$ gives a universal version of the connecting morphism. On the other hand, note that

$$\pi_1 \text{Hom}(T, K(\mathcal{O}, 3)) = H^2(T, \mathcal{O}) = \mathcal{O}$$

and

$$\pi_2 \text{Hom}(T, K(\mathcal{O}, 2)) = H^0(T, \mathcal{O}) = \mathcal{O}.$$

Thus the universal connecting morphism is a scalar constant C , and for any $\rho : Y \rightarrow T$ the connecting morphism for $\text{Hom}(Y, T)$ based at ρ fits into the diagram

$$\begin{array}{ccc} H^0(T, \mathcal{O}) & \rightarrow & H^2(T, \mathcal{O}) \\ \downarrow & & \downarrow \\ H^0(Y, \mathcal{O}) & \rightarrow & H^2(Y, \mathcal{O}). \end{array}$$

The vertical maps are those induced by ρ . It follows that the connecting map for $\text{Hom}(Y, T)$ based at ρ is the same constant C multiplied by the class η which is the

image of ρ in $H^2(Y, \mathcal{O})$. For the claim, apply this to $Y = X_{UDol}$. To finish proving the claim we just have to show that C is nonzero. But if C were zero then the generator for $\pi_2(Hom(T, K(\mathcal{O}, 2)), 1_T)$ would lift to an element of $\pi_2(Hom(T, T), 1_T)$. This would give a map

$$S^2 \times T \rightarrow T$$

(where S^2 denotes the constant presheaf with values S^2), which is nontrivial on $S^2 \times \{0\}$ and $\{0\} \times T$. This map would correspond to an element $\mu \in H^2(S^2 \times T, \mathcal{O})$ with $\mu \cup \mu = 0$. But we have

$$H^2(S^2 \times T, \mathcal{O}) = H^2(S^2, \mathcal{O}) \oplus H^2(T, \mathcal{O}) = \mathcal{O} \oplus \mathcal{O}$$

and the cup product of the two components is nontrivial (by Künneth). Therefore it is impossible to have a class μ which is nontrivial in both components but with $\mu \cup \mu = 0$. This contradiction implies that $C \neq 0$, giving the claim. $\square$

With the claim we obtain

$$\pi_1(R, \rho) \cong H_{Dol}^2(X)/(\eta).$$

Thus our characteristic class κ becomes a map

$$\kappa : \bigwedge^2(W) \rightarrow H_{Dol}^2(X)/(\eta).$$

Next, we choose η (which fixes the choice of ρ up to homotopy). Let E be the exceptional divisor on X and let H denote the pullback of an ample divisor on Z not meeting the point P . Let $[E]$ and $[H]$ denote their Chern classes in $H^1(X, \Omega_X^1) \subset H_{Dol}^2(X)$. We set

$$\eta = m[E] + n[H] \in H_{Dol}^2(X).$$

We have to choose n and m so that $\eta \cup \eta = 0$. Note that $H_{Dol}^4(X) \cong \mathbf{C}$ with natural morphism given by the residue map, normalized so that the cohomology class of a point is equal to 1. Via this isomorphism, $[E] \cup [E] = E.E = -1$ and $[H] \cup [H] = H.H \in \mathbf{Z}$. Note that $[E] \cup [H] = 0$ since the two divisors don't intersect. Thus

$$\eta \cup \eta = n^2 H.H - m^2.$$

We choose m, n so that this is equal to 0.

Remark: If $H.H$ is not the square of an integer, then the pair (m, n) can not be chosen in $\mathbf{Q}^2$, and in particular our map will not exist as a topological map $X^{\text{top}} \rightarrow S^2$. However, our map will exist as a map from the constant presheaf with values X^{top} , to $S^2 \otimes \mathbf{C}$. (This is a heuristic remark since in the present paper we don't treat the question of the relationship between Betti cohomology and Dolbeault cohomology).

Here is the result of the calculation which will be done below.

Theorem 7.1 *With the above choices of T , X , $R := \text{Hom}(X_{U_{\text{Dol}}}, T)$, $\eta = m[E] + n[H]$ ($\eta \neq 0$), and hence $\rho \in R(\text{Spec } \mathbf{C})$, the secondary Kodaira-Spencer class*

$$\kappa : \bigwedge^2 W \rightarrow \pi_1(R, \rho) = H_{\text{Dol}}^2(X)/(\eta)$$

lands in $H^2(X, \mathcal{O}) \subset H_{\text{Dol}}^2(X)/(\eta)$ and the map

$$\bigwedge^2 W \rightarrow H^2(X, \mathcal{O})$$

is dual (using Serre duality, the isomorphism $H^0(X, \Omega_X^2) \cong H^0(Z, \Omega_Z^2)$, and the isomorphism $W^ \cong T^*(Z)_P$) to $m^2 \text{ev}_P$ where*

$$\text{ev}_P : H^0(Z, \Omega_Z^2) \rightarrow \bigwedge^2 T^*(Z)_P$$

is the evaluation morphism. In particular, $\kappa \neq 0$ if $h^{2,0}(X) = h^{2,0}(Z) > 0$ and $m \neq 0$.

Before getting on with the proof in §8 below, we make a few general remarks about this result. A similar thing can be stated for the “Dolbeault homotopy type of X ”. One way of defining this (which wouldn’t be the historical way, though) is as the 1-connected very presentable n -stack Σ representing the very presentable shape of X_{Dol} (cf Theorem 10.1 in Appendix II below). In this point of view, we get an action of W on the Dolbeault homotopy type. The theorem says that this action of W is nontrivial. Note however that the action of W on the homotopy group sheaves (which are the homotopy groups of X tensored with $\mathbf{C}$) will be trivial. It is certainly possible to define the action of W , and to make the same calculation as below to show that the action is nontrivial, using the algebra of forms

$$A_{\text{Dol}}(X) = \left(\bigoplus_{p,q} A^{p,q}(X), \bar{\partial} \right).$$

In fact, my first heuristic version of the calculation was done using forms. However, the technical details relating a differential-forms version of nonabelian cohomology, with the version presented here, seem for the moment somewhat difficult, so we restrict in the present paper to an algebraic version of the calculation.

The secondary class κ is a natural map, so it doesn’t really have any choice other than to be a multiple of the dual of the evaluation map ev_P . The only question is whether this multiple is nonzero or not. Here is a heuristic global argument to see why, in principle, the constant should be nonzero. Let $\mathcal{X} \rightarrow Z$ be the total space of the family of blow-ups of points moving in Z . It is obtained by blowing up the diagonal in $Z \times Z$. The secondary Kodaira-Spencer class we calculate here is (or should be, at least) the $(2, 0) \times (0, 2)$ Hodge component (i.e. the component of type $(2, 0)$ on the base and $(0, 2)$ on the fiber) of the

following globally defined invariant. Fix $\eta \in H^2(X, \mathbf{C})$, which is invariant under the monodromy since the monodromy is trivial (the base Z being simply connected); the degeneration of the Leray spectral sequence says that η comes from restriction of a global class $\tilde{\eta} \in H^2(\mathcal{X}, \mathbf{C})$; the cup product $\tilde{\eta} \cup \tilde{\eta}$ restricts to zero on the fibers, so it lies in the next step for the Leray filtration, which in our case is $H^2(Z, H^2(X, \mathbf{C}))$. This cup product is therefore a globally defined class. It is the obstruction to extending $\rho : X^{\text{top}} \rightarrow T$ to a map $\mathcal{X}^{\text{top}} \rightarrow T$. The $(2, 0) \times (0, 2)$ component of this class, which is a holomorphic 2-form on Z with coefficients in $H^2(X, \mathcal{O})$, *should* give κ when evaluated at the point $P \in Z$. I don't currently have a proof of this, though.

In our example, it is relatively easy to see by looking at the cohomology class of the diagonal that the $(2, 0) \times (0, 2)$ component of the global class is nonzero, and in fact it is the identity matrix (via Serre duality). Thus if one could prove the above statement that the global class gives c under evaluation at $P \in Z$, then this would prove that $\kappa \neq 0$ when $H^2(X, \mathcal{O}) \neq 0$.

With the previous paragraphs as heuristic argument, the result that $\kappa \neq 0$ doesn't look all that surprising. Still, it means that the “variation of nonabelian Hodge structure”² on the family of $\text{Hom}(X^{\text{top}}, T)$, when X is a variable fiber in the family $\mathcal{X} \rightarrow Z$, is nontrivial, and even nontrivial for infinitesimal reasons. The base of this “variation” is Z which is simply connected. In particular, the variations of mixed Hodge structure on the homotopy groups (or anything else you could think of) are trivial. From a topological point of view, it is never a surprise to find a family where the homotopy groups are constant but the family nontrivial. On the other hand, this goes against the commonly held intuition for projective algebraic varieties that “formality means that everything is determined by the cohomology ring”: in the example $\mathcal{X} \rightarrow Z$, the locally constant family—parametrized by Z^{top} —of cohomology rings of the fibers X is trivial. What remains true of course is that the topology of the family is determined by the cohomology ring of the total space $\mathcal{X}$. Our secondary Kodaira-Spencer invariant is a local invariant which contributes to nontriviality of the global cohomology ring of the total space of the family.

Our class detects the motion of a point $P \in Z$ exactly when $H^0(Z, \Omega_Z^2) \neq 0$. This seems to fit in with the standard intuition that $H^0(Z, \Omega_Z^2) \neq 0$ causes the class group of zero cycles to be big (Mumford's and Clemens' results, Bloch conjecture etc. cf Voisin [48]). I don't see a precise connection, though.

8. The calculation

We keep the above notations $T, P \in Z, X, E, H, \eta, \rho$. We establish some more: let N be an affine neighborhood of P in Z such that TZ is trivialized over N . Assume

² This terminology is put in quotes because the current discussion of Dolbeault cohomology is only a first step towards defining what a “variation of nonabelian Hodge structure” is.

that N doesn't meet the divisor image of H in Z . Let α and β denote basis sections in $TZ(N)$. Let B be the inverse image of N in X , and let $C = X - E \cong Z - \{P\}$. Then put $A := B \cap C$. Note that $\{B, C\}$ is an open covering of X . We can write

$$X = B \cup^A C$$

(this is true as a pushout of sheaves of sets on $Sch/\mathbf{C}$). Similarly we have

$$X_{UDol} = B_{UDol} \cup^{A_{UDol}} C_{UDol}.$$

The basis vectors α, β give cocycles for elements in $H^1(X, TX)$ (actually in the Čech cohomology relative to our covering) and it is easy to see that these cocycles project to basis elements of our 2-dimensional space W . We denote the basis vectors of W also by α and β .

We can describe the action of W on X_{UDol} concretely in the following way. Set $K := K(W, 1)$ with basepoint denoted $0 \in K$. Then the trivialization of $TZ|_N$ gives an equivalence

$$A_{UDol} \cong A \times K.$$

There is a group structure on K , that is a morphism $K \times K \rightarrow K$ corresponding to the addition on W , and this gives

$$A \times K \times K \rightarrow A \times K,$$

which we can rewrite as

$$\mu : A_{UDol} \times K \rightarrow A_{UDol}.$$

Putting the identity 1_K in the second variable we get a map

$$\Phi : A_{UDol} \times K \rightarrow A_{UDol} \times K$$

which is an equivalence. Note also that $\Phi|_{A_{UDol} \times 0}$ is the identity of A_{UDol} .

Let i, j be the inclusions from A_{UDol} to B_{UDol} and C_{UDol} respectively. Then use the inclusions

$$(i \times 1_K) \circ \Phi : A_{UDol} \times K \rightarrow B_{UDol} \times K$$

and

$$j \times 1_K : A_{UDol} \times K \rightarrow C_{UDol} \times K$$

to construct the pushout

$$P := B_{UDol} \times K \cup^{A_{UDol} \times K} C_{UDol} \times K.$$

This comes equipped with a morphism

$$P \rightarrow K$$

and the fiber over the basepoint $0 \in K$ is just

$$B_{UDol} \cup^{A_{UDol}} C_{UDol} = X_{UDol}.$$

Therefore according to our definition, $P \rightarrow K$ is an action of W on X_{UDol} .

The corresponding action of K on $R := \text{Hom}(X_{UDol}, T)$ is by definition

$$\text{Hom}(P/K, T) \rightarrow K.$$

Using the Mayer-Vietoris principle we get

$$\begin{aligned} \text{Hom}(P/K, T) = \\ \text{Hom}(B_{UDol} \times K/K, T) \times_{\text{Hom}(A_{UDol} \times K/K, T)} \text{Hom}(C_{UDol} \times K/K, T). \end{aligned}$$

However, note that

$$\text{Hom}(B_{UDol} \times K/K, T) = \text{Hom}(B_{UDol}, T) \times K$$

and similarly for the other factors. The morphism

$$\text{Hom}(C_{UDol}, T) \times K \rightarrow \text{Hom}(A_{UDol}, T) \times K$$

induced by $j \times 1_K$ is just the product of the morphism induced by j , with 1_K . (On the other hand, the same is not true of the first morphism in the fiber product, as it is induced by $(i \times 1_K) \circ \Phi$.) We can now write

$$\begin{aligned} \text{Hom}(P/K, T) = \\ (\text{Hom}(B_{UDol}, T) \times K) \times_{\text{Hom}(A_{UDol}, T)} \text{Hom}(C_{UDol}, T). \end{aligned}$$

The first morphism in the fiber product is the composition of the product-compatible morphism

$$(j^* \times 1_K) : \text{Hom}(B_{UDol}, T) \times K \rightarrow \text{Hom}(A_{UDol}, T) \times K,$$

with the morphism

$$\Psi : \text{Hom}(A_{UDol}, T) \times K \rightarrow \text{Hom}(A_{UDol}, T).$$

This map is equivalent (by the definition of internal Hom) to

$$\text{Hom}(A_{UDol}, T) \rightarrow \text{Hom}(K, \text{Hom}(A_{UDol}, T)),$$

which in turn is equivalent to a map

$$Hom(A_{UDol}, T) \rightarrow Hom(A_{UDol} \times K, T),$$

this latter being induced by our action $\mu : A_{UDol} \times K \rightarrow A_{UDol}$.

The first step is to notice that the map $\rho|_{A_{UDol}}$ from A_{UDol} to T factors through a map

$$h : A_{UDol} \rightarrow K(\mathcal{O}, 3) \rightarrow T.$$

This factorization is given by the fact that η is a class in $H^2((X_{Dol}, A_{Dol}), \mathcal{O})$, in other words we are given a trivialization of η over A_{Dol} .

Recall that we write $\rho = (\eta, \varphi)$ where $\eta : X_{UDol} \rightarrow K(\mathcal{O}, 2)$ and φ is a section of the pullback bundle

$$L_\eta := X_{UDol} \times_{K(\mathcal{O}, 4)} *$$

which is a bundle with fiber $K(\mathcal{O}, 3)$ over X_{UDol} . The section φ determines a trivialization

$$L_\eta \cong X_{UDol} \times K(\mathcal{O}, 3)$$

such that φ corresponds to the 0-section. This trivialization is uniquely determined by the condition that it be compatible with the structure of “principal bundle” under the “group” $K(\mathcal{O}, 3)$.

We adopt the following strategy for calculating the secondary class. We will look at new n -stacks $P^i \rightarrow K$ with K -maps $P^i \rightarrow P$, and points $\rho^i \in P_0^i$ (where P_0^i means the fiber of P^i over $0 \in K$), such that ρ^i maps to ρ . We have to arrange so that the primary class is trivial, in other words that the class of ρ^i in $\pi_0(P_0^i)$ should be invariant under the action of $W = \pi_1(K)$. We also have to insure that the other primary class, the action of W on $\pi_1(P_0^i, \rho^i)$ by outer automorphisms, should be trivial. In this case, we obtain a secondary class for ρ^i , which is an element of

$$H^2(K, \pi_1(P_0^i, \rho^i))$$

and this secondary class maps to our class for P .

First of all, let $*$ $\rightarrow Hom(B_{UDol}, T)$ be the morphism corresponding to the point $\rho|_{B_{UDol}}$. We get

$$K = * \times K \rightarrow Hom(B_{UDol}, T) \times K.$$

Thus we obtain a morphism

$$\begin{aligned} P^1 &:= K \times_{Hom(A_{UDol}, T)} Hom(C_{UDol}, T) \rightarrow \\ &(Hom(B_{UDol}, T) \times K) \times_{Hom(A_{UDol}, T)} Hom(C_{UDol}, T) = P. \end{aligned}$$

This morphism is compatible with the projections to K .

The first morphism

$$u : K \rightarrow \text{Hom}(A_{UDol}, T)$$

in the fiber product is obtained by the composition

$$\begin{aligned} K &\rightarrow \text{Hom}(B_{UDol}, T) \times K \xrightarrow{(i \times 1_K) \circ \Phi} \\ &\text{Hom}(A_{UDol}, T) \times K \xrightarrow{p_1} \text{Hom}(A_{UDol}, T). \end{aligned}$$

Thus u corresponds to the map $K \times A_{UDol} \rightarrow T$ obtained by composing

$$K \times A_{UDol} \xrightarrow{\mu} A_{UDol} \xrightarrow{\rho} T.$$

The map u factors through a morphism

$$K \rightarrow \Gamma(A_{UDol}, L_\eta) \rightarrow \text{Hom}(A_{UDol}, T)$$

(technically speaking what should enter into the above notation is $L_\eta|_{A_{UDol}}$ but for brevity we omit the restriction since it is implicitly determined by the notation $\Gamma(A_{UDol}, -)$.) The factorization comes about because the composition

$$A_{UDol} \xrightarrow{\rho} T \rightarrow K(\mathcal{O}, 2),$$

is given as the constant map at the basepoint. Thus pulling back by

$$A_{UDol} \times K \rightarrow A_{UDol} \rightarrow K(\mathcal{O}, 2)$$

is again constant at the basepoint so the map

$$K \rightarrow \text{Hom}(A_{UDol}, T)$$

factors through a map

$$\tilde{u} : K \rightarrow \text{Hom}(A_{UDol}, K(\mathcal{O}, 3)).$$

Following above, $\tilde{u}$ corresponds to the composition

$$K \times A_{UDol} \xrightarrow{\mu} A_{UDol} \xrightarrow{h} K(\mathcal{O}, 3),$$

which we can write as $\tilde{u} = \mu^*(h)$.

Now we obtain a morphism (over K)

$$P^2 := K \times_{\Gamma(A_{UDol}, L_\eta)} \Gamma(C_{UDol}, L_\eta)$$

$$\rightarrow K \times_{\text{Hom}(A_{UDol}, T)} \text{Hom}(C_{UDol}, T) = P^1.$$

We have used the section φ of L_η to obtain a trivialization

$$L_\eta \cong X_{UDol} \times K(\mathcal{O}, 3).$$

Via this equivalence the section corresponding to ρ (that is, the section φ) corresponds to the zero-section. Using the trivialization given by φ we can write

$$\Gamma(A_{UDol}, L_\eta) \cong \text{Hom}(A_{UDol}, K(\mathcal{O}, 3))$$

(*Caution:* this is not the same trivialization as given by saying that A_{UDol} maps to the basepoint of $K(\mathcal{O}, 2)$ so the pullback fibration L_η is trivial over A_{UDol} ; these two trivializations differ by translation by h , a point which will come up below); and

$$\Gamma(C_{UDol}, L_\eta) \cong \text{Hom}(C_{UDol}, K(\mathcal{O}, 3)).$$

These are compatible with the restriction j^* so we can write

$$P^2 = K \times_{\text{Hom}(A_{UDol}, K(\mathcal{O}, 3))} \text{Hom}(C_{UDol}, K(\mathcal{O}, 3)),$$

with the second morphism in the fiber product being the restriction j^* acting on maps to $K(\mathcal{O}, 3)$.

We have to re-calculate the first morphism in the fiber product

$$a : K \rightarrow \text{Hom}(A_{UDol}, K(\mathcal{O}, 3)).$$

It is no longer equal to $\tilde{u} = \mu^*(h)$, because when we set the section φ of L_η equal to the zero-section to get an equivalence between L_η and $K(\mathcal{O}, 3)$, this made a translation on $L_\eta|_{A_{UDol}}$ —which was already trivial due to the fact that $\eta|_{A_{UDol}} = 0$ —this translation has the effect of setting h equal to the 0-section. This translation gives us the formula

$$a = \mu^*(h) - p_2^*(h)$$

where $p_2^*(h)$ is the map $K \rightarrow \text{Hom}(A_{UDol}, K(\mathcal{O}, 3))$ corresponding to the composition

$$K \times A_{UDol} \xrightarrow{p_2} A_{UDol} \xrightarrow{h} K(\mathcal{O}, 3).$$

Note that p_2 denotes the second projection. The minus sign in the equation for a is subtraction using the “abelian group” (i.e. E_∞) structure of $\text{Hom}(A_{UDol}, K(\mathcal{O}, 3))$ induced by the “abelian group” structure of $K(\mathcal{O}, 3)$.

We now turn back to the second morphism in the fiber product,

$$j^* : \text{Hom}(C_{UDol}, K(\mathcal{O}, 3)) \rightarrow \text{Hom}(A_{UDol}, K(\mathcal{O}, 3)).$$

The infinite loop space structure of $K(\mathcal{O}, 3)$ which is inherited by $Hom(\mathcal{F}, K(\mathcal{O}, 3))$ for $\mathcal{F} = C_{UDol}$ and $\mathcal{F} = A_{UDol}$. By Proposition 1.1, this delooping structure gives a decomposition of $Hom(\mathcal{F}, K(\mathcal{O}, 3))$ into a product of Eilenberg-MacLane stacks. The restriction morphism j^* above is compatible with the delooping structures, so by Proposition 1.1 (B), it is homotopic to a map compatible with the decomposition into a product of Eilenberg-MacLane stacks.

Recall that

$$\pi_i(Hom(C_{UDol}, K(\mathcal{O}, 3)), 0) = H^{3-i}(C_{UDol}, \mathcal{O}) = H_{Dol}^{3-i}(C)$$

and

$$\pi_i(Hom(A_{UDol}, K(\mathcal{O}, 3)), 0) = H^{3-i}(A_{UDol}, \mathcal{O}) = H_{Dol}^{3-i}(A).$$

From the splitting given by the delooping structures via Proposition 1.1 with homotopy of functoriality of part (B) of that proposition (choose one), we obtain a homotopy-commutative diagram

$$\begin{array}{ccc} K(H_{Dol}^2(C), 1) & \rightarrow & K(H_{Dol}^2(A), 1) \\ \downarrow & & \downarrow \\ Hom(C_{UDol}, K(\mathcal{O}, 3)) & \rightarrow & Hom(A_{UDol}, K(\mathcal{O}, 3)). \end{array}$$

All maps are infinite loop maps. Using this diagram, we get the map

$$\begin{aligned} P^3 &:= K \times_{Hom(A_{UDol}, K(\mathcal{O}, 3))} K(H_{Dol}^2(C), 1) \rightarrow \\ &K \times_{Hom(A_{UDol}, K(\mathcal{O}, 3))} Hom(C_{UDol}, K(\mathcal{O}, 3)) = P^2. \end{aligned}$$

Set $\rho^3 := (0, 0)$ in P_0^3 . It maps to ρ^2 (a point which we didn't specify because it was always obviously given by ρ), because we have normalized so that φ becomes the zero-section. The map $P^3 \rightarrow P^2$ (obviously a map over K) takes ρ^3 to ρ^2 .

We have

$$H_{Dol}^2(C) = H^2(C, \mathcal{O}_C) \oplus H^1(C, \Omega_C^1) \oplus H^0(C, \Omega_C^2),$$

and similarly

$$H_{Dol}^2(A) = H^2(A, \mathcal{O}_A) \oplus H^1(A, \Omega_A^1) \oplus H^0(A, \Omega_A^2).$$

Recall that C and A are isomorphic to open subsets of Z : together N and C form a covering of Z and $N \cap C = A$. On the other hand, cohomology of coherent sheaves on N vanishes because N is affine. Therefore Mayer-Vietoris gives an exact sequence

$$H^1(C, \Omega_C^1) \rightarrow H^1(A, \Omega_A^1) \rightarrow H^2(Z, \Omega_Z^1).$$

We are supposing that $H_{Dol}^1(Z) = 0$ so by duality $H_{Dol}^3(Z) = 0$. Thus the term on the right is zero, and the morphism of vector spaces

$$H^1(C, \Omega_C^1) \rightarrow H^1(A, \Omega_A^1)$$

is surjective. Choose a splitting. This gives a morphism

$$K(H^1(A, \Omega_A^1), 1) \rightarrow K(H_{Dol}^2(C), 1)$$

such that the projection into $K(H_{Dol}^2(A), 1)$ is equal to the inclusion from the Dolbeault direct sum decomposition for A . Using our choice of splitting we obtain a map

$$P^4 := K \times_{Hom(A_{UDol}, K(\mathcal{O}, 3))} K(H^1(A, \Omega_A^1), 1) \rightarrow$$

$$K \times_{Hom(A_{UDol}, K(\mathcal{O}, 3))} K(H_{Dol}^2(C), 1) = P^3.$$

Again set $\rho^4 = (0, 0) \in P_0^4$, which maps to ρ^3 .

Now P^4 is defined entirely in terms of A . We decompose things a bit more. Using the infinite loop-space structure of $Hom(A_{UDol}, K(\mathcal{O}, 3))$ and again Proposition 1.1 we get the decomposition into a product of Eilenberg-MacLane spaces

$$Hom(A_{UDol}, K(\mathcal{O}, 3)) \cong J^0 \times J^1 \times J^2 \times J^3$$

where

$$J^i = K(H_{Dol}^{3-i}(A), i).$$

The second map in the fiber product defining P^4 is homotopic (choose a homotopy) to one which is compatible with this decomposition, coming from the morphism

$$K(H^1(A, \Omega_A^1), 1) \rightarrow J^1 = K(H_{Dol}^2(A), 1)$$

(the other components are the maps sending $*$ to the basepoints $0 \in J^i$, $i \neq 2$).

Thus we can write

$$P^4 = Q^0 \times Q^1 \times Q^2 \times Q^3$$

where

$$Q^i = K \times_{J^i} * = K \times_{K(H_{Dol}^{3-i}(A), i)} *$$

for $i \neq 1$ and where

$$Q^1 = K \times_{K(H_{Dol}^2(A), 1)} K(H^1(A, \Omega_A^1), 1).$$

This means that the action of W on P_0^4 decomposes into a product of actions on the Q_0^i . Note that the Q_0^i are themselves Eilenberg-MacLane spaces,

$$Q_0^i = \Omega J^i = K(H_{Dol}^{3-i}(A), i-1)$$

for $i \geq 2$, and

$$Q_0^1 = K(H_{Dol}^2(A)/H^1(A, \Omega_A^1), 0).$$

The classifying maps for the actions of W on the components Q_0^i factor as

$$K \rightarrow J^i \rightarrow B \operatorname{Aut}(Q_0^i)$$

for $i \geq 2$, and

$$K \rightarrow K(H_{Dol}^2(A)/H^1(A, \Omega_A^1), 1) \rightarrow B \operatorname{Aut}(Q_0^1)$$

for $i = 1$. Thus the classifying map for the action $K \rightarrow B \operatorname{Aut}(P_0^4)$ factors through our above map

$$a : K \rightarrow K(H_{Dol}^2(A)/H^1(A, \Omega_A^1), 1) \times J^2 \times J^3.$$

We use this to calculate the characteristic classes for the action: the primary invariant is the first component, corresponding to a map

$$W \rightarrow H_{Dol}^2(A)/H^1(A, \Omega_A^1).$$

The other primary invariant giving the action of W on π_1 by outer automorphisms, is trivial: it is the action on ΩJ^2 induced by the classifying map $K \rightarrow J^2$, and J^2 is simply connected. The secondary invariant which we are interested in is the map $K \rightarrow J^2$ which corresponds to a class in $H^2(K, H_{Dol}^1(A))$.

A preliminary remark is that the formula $a = \mu^*(h) - p_2^*(h)$ means that the component of a in J^0 is equal to 0. In fact, $p_2^*(h)$ is exactly the J^0 -component of $\mu^*(h)$. On the other hand, the remaining components of a are the same as those of $\mu^*(h)$; these are the Künneth components of

$$h \circ \mu : K \times A_{UDol} \rightarrow K(\mathcal{O}, 3).$$

The first thing to check is that the primary invariant is trivial for (P^4, ρ^4) . As we have said above, it is the map

$$K(W, 1) \rightarrow K(H_{Dol}^2(A)/H^1(A, \Omega_A^1), 1).$$

To check that it is trivial, it suffices to prove the

Claim $(*)$: the map

$$W \rightarrow \pi_1(\operatorname{Hom}(A_{UDol}, K(\mathcal{O}, 3)), 0) = H_{Dol}^2(A)$$

takes W into the component $H^1(A, \Omega_A^1)$. For this we must again get back to the description of the map

$$\mu^*(h) : K \rightarrow \operatorname{Hom}(A_{UDol}, K(\mathcal{O}, 3))$$

(note as remarked above that all of the components except the J^0 component, are the same for $\mu^*(h)$ or for a). This map, which is equivalent to a map

$$A_{UDol} \times K \rightarrow K(\mathcal{O}, 3),$$

is the pullback of

$$h : A_{UDol} \rightarrow K(\mathcal{O}, 3)$$

by the map

$$\mu : A_{UDol} \times K \rightarrow A_{UDol}.$$

Now we have

$$A_{UDol} = A \times K$$

so

$$H^3(A_{UDol}, \mathcal{O}) = H^3(A, \mathcal{O}) \oplus H^2(A, H^1(K, \mathcal{O})) \oplus H^1(A, H^2(K, \mathcal{O})) \oplus H^3(K, \mathcal{O}).$$

The last term $H^3(K, \mathcal{O})$ vanishes because it would be $\bigwedge^3 W^*$ but W is 2-dimensional. Similarly, one can arrange (by an appropriate choice of N) so that A has an open covering by two affine open sets. Thus $H^3(A, \mathcal{O}) = H^2(A, \mathcal{O}) = 0$. The only remaining term is

$$H^3(A_{UDol}, \mathcal{O}) = H^1(A, H^2(K, \mathcal{O})) = H^1(A, \bigwedge^2 W^*) = H^1(A, \Omega_A^2).$$

Our map

$$A_{UDol} \times K \rightarrow K(\mathcal{O}, 3)$$

is obtained by pulling back the above, using the map $K \times K \rightarrow K$. Pullback for this map is

$$\begin{aligned} & \bigwedge^2 W^* \rightarrow \\ & H^2(K, \mathcal{O}) \otimes_{\mathcal{O}} \mathcal{O} \oplus H^1(K, \mathcal{O}) \otimes_{\mathcal{O}} H^1(K, \mathcal{O}) \oplus \mathcal{O} \otimes_{\mathcal{O}} H^2(K, \mathcal{O}). \end{aligned}$$

Each of the factors is nontrivial, with the middle being (up to a multiple which depends on normalizations for notation in exterior products) the standard map

$$\bigwedge^2 W^* \rightarrow W^* \otimes_{\mathcal{O}} W^*.$$

The morphism induced by pulling back h to $A_{UDol} \times K$ thus decomposes into Künneth components

$$H^1(A, \Omega_A^2) \otimes_{\mathcal{O}} \mathcal{O} \oplus H^1(A, \Omega_A^1) \otimes_{\mathcal{O}} W^* \oplus H^1(A, \mathcal{O}_A) \otimes_{\mathcal{O}} \bigwedge^2 W^*.$$

Each component is induced by $h \in H^1(A, \Omega_A^2)$.

The map

$$\pi_1(K) = W \rightarrow \pi_1(\text{Hom}(A_{UDol}, K(\mathcal{O}, 3))) = H_{Dol}^2(A)$$

corresponds to the middle component, which is in fact a map

$$W \rightarrow H^1(A, \Omega_A^1).$$

This was exactly the claim (*) we needed to prove to show that the primary invariant for (P^4, ρ^4) was trivial.

The other primary invariant, the action of W on $\pi_1(P_0^4, \rho^4)$, is trivial as remarked above. To restate the argument, the map a' induces an injection on π_1 , therefore the π_1 of the fiber is the image of π_2 of the base; but since the base is an infinite loop space, the action of π_1 of the base on π_2 of the base is trivial; thus the action of π_1 of the base and in particular of W on $\pi_1(P_0^4, \rho^4)$ is trivial.

We can now look at the secondary invariant for (P^4, ρ^4) . It is the map

$$K \rightarrow J^2 = K(H_{Dol}^1(A), 2),$$

which is the next Künneth component of the pullback of h to $A_{UDol} \times K$,

$$\bigwedge^2 W \rightarrow H^1(A, \mathcal{O}_A) \subset H_{Dol}^1(A) = \pi_2(\text{Hom}(A_{UDol}, K(\mathcal{O}, 3))).$$

This claim tells us that the secondary class for P^4 is just h considered as an element of

$$H^1(A, \Omega_A^2) = H^1(A, \bigwedge^2 W^*) = H^1(A, \mathcal{O}_A) \otimes_{\mathcal{O}} \bigwedge^2 W^*.$$

Recalling that

$$P_0^4 = * \times_{\text{Hom}(A_{UDol}, K(\mathcal{O}, 3))} K(H^1(A, \mathcal{O}_A), 1)$$

and $\rho^4 = (0, 0)$, we have that P_0^4 is just the homotopy fiber of the second morphism. The long exact sequence for the fibration gives

$$0 \rightarrow \pi_2(\text{Hom}(A_{UDol}, K(\mathcal{O}, 3))) \rightarrow \pi_1(P_0^4, \rho^4) \rightarrow 0$$

(the morphism from π_1 of the total space to π_1 of the base being injective, and π_2 of the total space being zero). This long exact sequence persists under the maps

$$P_0^4 \rightarrow P_0^3 \rightarrow P_0^2 \rightarrow P_0^1$$

and furthermore, even into P_0 where the long exact sequence for the fiber of a morphism is replaced by a long exact sequence for the homotopy fiber product. It follows that the secondary class for (P, ρ) is the image of our above class (basically h)

$$\bigwedge^2 W \rightarrow H^1(A, \mathcal{O}_A)$$

under composition with the map

$$\begin{aligned} H^1(A, \mathcal{O}_A) &= \pi_2(\text{Hom}(A_{UDol}, K(\mathcal{O}, 3)), 0) \rightarrow \pi_2(\text{Hom}(A_{UDol}, T), \rho|_{A_{UDol}}) \\ &\rightarrow \pi_1(\text{Hom}(X_{UDol}, T), \rho) = H_{Dol}^2(X)/(\eta). \end{aligned}$$

One slight twist to notice is that the morphism $K(\mathcal{O}, 3) \rightarrow T$ in question (over A_{UDol}) is shifted by h . This shift is recovered in the last equality, where we undo a shift by φ .

This morphism from the long exact sequence for the fiber product (of Hom 's) is equal to the connecting morphism

$$H^1(A, \mathcal{O}_A) \rightarrow H^2(X, \mathcal{O}_X) \subset H_{Dol}^2(X)/(\eta).$$

Finally, we have concluded that our secondary class is the composition

$$\bigwedge^2 W^* \rightarrow H^1(A, \mathcal{O}_A) \rightarrow H^2(X, \mathcal{O}_X)$$

where the first map is h (which depends on η and which we investigate below) and the second map is the connecting morphism.

The remaining problem is to calculate $h \in H^1(A, \Omega_A^2)$ or more precisely its image by the connecting morphism. We have the exact sequence of the cohomology of the pair (X, A) with coefficients in Ω_X^2 :

$$\begin{aligned} 0 &= H^1(X, \Omega_X^2) \rightarrow H^1(A, \Omega_A^2) \rightarrow \\ &H^2((X, A), \Omega_X^2) \rightarrow H^2(X, \Omega_X^2). \end{aligned}$$

The class η may be considered as lying in $H^1((X, A), \Omega_X^1)$, which is the statement that our map to T factors, over A , through a map to $K(\mathcal{O}, 3)$. Therefore the cup product $\eta \cup \eta$ can be considered as lying in $H^2((X, A), \Omega_X^2)$ and mapping to zero in $H^2(X, \Omega_X^2)$. The class $h \in H^1(A, \Omega_A^2)$ is the preimage of $\eta \cup \eta$ in the exact sequence for the pair (X, A) .

Now recall that $X = B \cup C$. This means that the pair (X, A) decomposes as a “disjoint union” of the pairs (X, B) and (X, C) (after applying excision). Thus we can write

$$H^2((X, A), \Omega_X^2) = H^2((X, B), \Omega_X^2) \oplus H^2((X, C), \Omega_X^2).$$

Our class $h \in H^1(A, \Omega_A^2)$ corresponds to $b + c$ with

$$b = -n^2[H]^2 \in H^2((X, B), \Omega_X^2)$$

and

$$c = m^2[E]^2 \in H^2((X, C), \Omega_X^2).$$

We now do the same thing for $Z = N \cup C$. The class h corresponds here to an element of

$$H^2((Z, N), \Omega_Z^2) \oplus H^2((Z, C), \Omega_Z^2).$$

This is again of the form $b' + c'$ but now with

$$b' = -n^2[H]^2 \in H^2((Z, N), \Omega_Z^2)$$

and

$$c' = m^2[P] \in H^2((Z, C), \Omega_Z^2).$$

From the long exact sequence of the pair (Z, N) and the fact that N is affine we find that

$$H^2((Z, N), \Omega_Z^2) \cong H^2(Z, \Omega_Z^2) = \mathbf{C}.$$

To calculate our secondary class we have to contract h with $\alpha \wedge \beta$ to obtain a class in $H^1(A, \mathcal{O}_A)$ and then take its image by the connecting map in $H^2(X, \mathcal{O}_X)$ or equivalently in $H^2(Z, \mathcal{O}_Z)$. To measure this image, use Serre duality: we will choose a form $\omega \in H^0(Z, \Omega_Z^2)$ and take the cup-product to end up with a class in $H^2(Z, \Omega_Z^2)$ (of which we then take the residue to end up in $\mathbf{C}$). The result of this procedure is the same as if we first contract ω with $\alpha \wedge \beta$ and then multiply this section of $H^0(A, \mathcal{O}_A)$ by h getting a class in $H^1(A, \Omega_A^2)$. Then take the image of this class by the connecting map and take its residue. Note that the contraction of ω with $\alpha \wedge \beta$ is defined over all of the neighborhood N . Call this section $\lambda \in \mathcal{O}(N)$. We are now reduced to the problem of calculating the image in $H^2(Z, \Omega_Z^2)$ under the connecting map for the covering $Z = N \cup C$, of $\lambda h \in H^1(A, \Omega_A^2)$.

We have written that the image of h in $H^2((Z, A), \Omega_Z^2)$ (which we shall denote $[h]$) is equal to $b' + c'$ where $b' \in H^2((Z, N), \Omega_Z^2)$ and $c' \in H^2((Z, C), \Omega_Z^2)$. The components b' and c' are obtained by residue maps for the class h , along respectively H and P (noting that $H = Z - N$ and $P = Z - C$). The form of the residue map is not important for us, just the fact that the classes have poles of order 1; it follows that if λ is a regular function on N (a neighborhood of P) then the residue of λh at P , is equal to λ times the residue of h at P . Therefore we can write

$$[\lambda h] = b'' + \lambda c',$$

with

$$b'' \in H^2((Z, N), \Omega_Z^2)$$

and

$$\lambda c' \in H^2((Z, C), \Omega_Z^2)$$

both being obtained by excision.

The value of $b'' \in H^2((Z, N), \Omega_Z^2) \cong \mathbf{C}$ is determined by the condition that the image of $[\lambda h]$ in $H^2(Z, \Omega_Z^2)$ be zero (from the long exact sequence for the pair (Z, A)).

We now look at the image of λh by the connecting map in the long exact sequence of the covering $Z = N \cup C$,

$$\begin{aligned} H^1(N, \Omega_N^2) \oplus H^1(C, \Omega_C^2) &\rightarrow H^1(A, \Omega_A^2) \\ &\rightarrow H^2(Z, \Omega_Z^2) \rightarrow \dots \end{aligned}$$

We can decompose this connecting map as a composition

$$H^1(A, \Omega_A^2) \rightarrow H^2((Z, A), \Omega_Z^2) \rightarrow H^2((Z, C), \Omega_Z^2) \rightarrow H^2(Z, \Omega_Z^2),$$

where the second arrow is the projection onto the first factor in the excision decomposition

$$H^2((Z, A), \Omega_Z^2) = H^2((Z, C), \Omega_Z^2) \oplus H^2((Z, N), \Omega_Z^2)$$

One could equally well use the second factor, with a sign change; our calculations are not accurate insofar as signs are concerned.

Thus the image of λh by the connecting map for the covering $Z = N \cup C$, is equal to the class of either $\lambda c'$ or of $-b''$. We don't know how to calculate b'' so we use the representation as $\lambda c'$. This image is then equal to

$$\lambda \cdot (m^2[P])$$

which is just $m^2\lambda(P)$ (because as noted above, $[P]$ is represented by cocycles with poles of order 1).

We have established the formula that the image of $\alpha \wedge \beta$ under the map

$$\bigwedge^2 W \rightarrow H^2(Z, \mathcal{O}_Z)$$

is a class which, when paired with a form $\omega \in H^0(Z, \Omega_Z^2)$, gives

$$m^2\omega(\alpha \wedge \beta)(P).$$

This completes the proof of Theorem 7.1. □

9. APPENDIX I: Relative Breen calculations in characteristic 0

Crucial to the reasonable working of a theory of nonabelian cohomology is the possibility of calculating the invariants in the Postnikov tower of the spaces which measure the

“shape”. In our setup, this means that we would like to calculate $H^i(K(\mathcal{O}, m), \mathcal{O})$. This calculation is the algebraic analogue of the classical Eilenberg-MacLane calculations. The algebraic version is the subject of Breen’s work [8], [9]. His motivation came mostly from arithmetic geometry, so he concentrated on the case of characteristic p in [9]. The characteristic 0 version, while not explicitly stated in [9], is implicit there because it is strictly easier than the characteristic p case: there are no new classes coming from Frobenius.

These calculations for the case of base scheme $\text{Spec}(\mathbf{C})$ are sufficient for the purposes of the present paper, but eventually a relative version will also be useful. In the context of calculation of Ext sheaves (i.e. the stable part of the calculation) this relative version was already evoked in [8], where Breen states that the $\text{Ext}^i(G, \cdot)$ for representable group schemes G can always be calculated. This part of the topic was not really taken up afterward, probably for lack of a reasonable category of sheaves over a base scheme S .

Such a category of sheaves is provided by Hirschowitz’s notion of *U-coherent sheaf* [21], see also Jaffe’s recent paper [26].³ We change Hirschowitz’s notation and call these objects *vector sheaves*. The category of vector sheaves over a base S is defined in [21] as the smallest abelian subcategory of sheaves of $\mathcal{O}$ -modules on the big site Sch/S , containing $\mathcal{O}$ and stable under localization of the base. Thus, locally over S vector sheaves are obtained starting with $\mathcal{O}$ by repeated applications of taking direct sums, kernels and cokernels. The abelian category of vector sheaves has several nice properties [21]. The coherent sheaves on S , which are defined as cokernels

$$\mathcal{O}^a \rightarrow \mathcal{O}^b \rightarrow \mathcal{F} \rightarrow 0,$$

are vector sheaves. Coherent sheaves are injective objects. Their duals, which we call *vector schemes*, are the group-schemes with vector space structure (but not necessarily flat) over S . These admit dual presentations as kernels of maps $\mathcal{O}^b \rightarrow \mathcal{O}^a$. They are projective objects (at least if the base S is affine). If S is affine, then any vector sheaf U admits resolutions

$$0 \rightarrow V \rightarrow V' \rightarrow V'' \rightarrow U \rightarrow 0$$

with V , V' and V'' vector schemes; and

$$0 \rightarrow U \rightarrow \mathcal{F} \rightarrow \mathcal{F}' \rightarrow \mathcal{F}'' \rightarrow 0$$

with $\mathcal{F}$, $\mathcal{F}'$, $\mathcal{F}''$ coherent sheaves.

³ Hirschowitz’s notion is similar to, but not quite the same as Auslander’s theory of “coherent functors” developed in the 1960’s. Jaffe’s paper [26] views *U-coherent sheaves* as a modification or generalization of Auslander’s theory—one looks at functors of algebras rather than functors of modules. Jaffe, who seems to have been unaware of Hirschowitz’s paper, cites an unpublished letter from Artin to Grothendieck, dating from the 1960’s, as a reference for the generalized version of Auslander’s theory. In order to straighten out the history of this notion, one would have to compare Artin’s letter with [21].

Example: The motivating example for the definition of vector sheaf in [21] was the following example. If E is a complex of vector bundles over S then the cohomology sheaves defined on the big site Sch/S are vector sheaves. Indeed they are vector sheaves of a special type which Hirschowitz calls “cohomologies”: quotients of vector schemes by coherent sheaves. This example is important because it arises from the cohomology of flat families of coherent sheaves parametrized by S : if $f : X \rightarrow S$ is a projective morphism and $\mathcal{F}$ is a coherent sheaf on X flat over S then a classical result says that the higher direct image complex $Rf_*(\mathcal{F})$ (calculated on the big site) is quasiisomorphic to a complex of vector bundles. The notion of vector sheaf (“ U -coherent sheaf” in [21]) thus keeps track of the jumping of cohomology of flat families of coherent sheaves.⁴

The most surprising property from [21] is that the duality functor $U^* := Hom(U, \mathcal{O})$ is exact, and is an involution. This is due to the fact that we take the big site Sch/S rather than the small Zariski or etale sites.

Another interesting point is that there are two different types of tensor products of vector sheaves: the *tensor product*

$$U \otimes_{\mathcal{O}} V := Hom(U, V^*)^*,$$

and the *cotensor product*

$$U \otimes^{\mathcal{O}} V := Hom(U^*, V).$$

These are not the same (although they coincide for coherent sheaves cf Lemma 9.1 below) and in particular they don’t have the same exactness properties. Neither of them is equal to the tensor product of sheaves of $\mathcal{O}$ -modules. See [39] and [41] for further discussion.

The above facts work in any characteristic and depend on the $\mathcal{O}$ -module structure. However, in characteristic zero vector sheaves have the additional property that the morphisms $U \rightarrow V$ of sheaves of abelian groups over $Sch/\mathbf{C}$ are automatically morphisms of

⁴ This type of example comes up in relation with Dolbeault cohomology: for example let

$$M := \mathcal{M}_{Dol}(X, G) = Hom(X_{Dol}, K(G, 1))$$

be the moduli stack of principal Higgs G -bundles. If V is a representation of G then we obtain

$$T := K(V/G, n) \rightarrow K(G, 1)$$

with fiber $K(V, n)$. There is a universal local system E on $X_{Dol} \times M$, and

$$\pi_i(Hom(X_{Dol}, T)/M, 0) = H^i(X_{Dol} \times M/M, E).$$

This is calculated by a Dolbeault complex for E on X/M , and the general discussion of cohomology in flat families applies. Therefore the $H^i(X_{Dol} \times M/M, E)$ are vector sheaves over M (here M is an algebraic stack; the condition of being a vector sheaf means that the pullback to any scheme $Y \rightarrow M$ is a vector sheaf on Y).

$\mathcal{O}$ -modules, see [39] [41]. Similarly, extensions of sheaves of abelian groups, between two vector sheaves, are again vector sheaves. These properties persist for the higher Ext^i , see Corollary 9.6 below. These properties do not remain true in characteristic p , as shown precisely by Breen's calculations of [9]. The basic problem is that Frobenius provides a morphism $\mathcal{O} \rightarrow \mathcal{O}$ of sheaves of abelian groups, which is not a morphism of sheaves of $\mathcal{O}$ -modules. This difficulty in characteristic p seems to be the main obstacle to realizing a reasonable analogue of rational homotopy theory, for homotopy in characteristic p . So we stick to characteristic 0!

We don't give a detailed introduction to vector sheaves, rather we refer the reader to [21], [39] and [41]. However, we do take this opportunity to correct an omission from [39] and [41]. Without the following lemma, the discussion in those references often seems contradictory, as tensor products and cotensor products are interchanged when the coefficients are coherent sheaves. For example, in the statement of Corollary 3.9 of [39] (which we restate as Theorem 9.2 and prove in more detail below), all terms occurring are coherent sheaves. Thus the tensor product which appears in the notation is also equal to the cotensor product. It is the cotensor product which appears most naturally in that situation. Indeed, Lemma 9.1 below explains (i.e. justifies) the seemingly erroneous statement $\mathcal{F} \otimes_{\mathcal{O}} \mathcal{G} = \underline{Hom}(\mathcal{F}^*, \mathcal{G})$ in the proof of Corollary 3.9 of [39].

Lemma 9.1 *Suppose $\mathcal{F}$ and $\mathcal{G}$ are coherent sheaves. Then the tensor product $\mathcal{F} \otimes_{\mathcal{O}} \mathcal{G}$ and the cotensor product $\mathcal{F} \otimes^{\mathcal{O}} \mathcal{G}$ coincide.*

Proof: Choose a presentation

$$\mathcal{O}^a \rightarrow \mathcal{O}^b \rightarrow \mathcal{G} \rightarrow 0.$$

Now $\mathcal{F}^* := Hom(\mathcal{F}, \mathcal{O})$ is a vector scheme, in particular it is a scheme affine over S . Therefore the functor

$$U \mapsto Hom(\mathcal{F}^*, U)$$

is exact in U . Applying this functor to the above presentation we obtain

$$\mathcal{F}^a \rightarrow \mathcal{F}^b \rightarrow Hom(\mathcal{F}^*, \mathcal{G}) \rightarrow 0.$$

The term $Hom(\mathcal{F}^*, \mathcal{G})$ is by definition the cotensor product $\mathcal{F} \otimes^{\mathcal{O}} \mathcal{G}$.

Taking the dual of the above presentation we obtain

$$0 \rightarrow \mathcal{G}^* \rightarrow \mathcal{O}^b \rightarrow \mathcal{O}^a.$$

Applying the functor

$$U \mapsto Hom(\mathcal{F}, U)$$

which is exact on the left, we obtain

$$0 \rightarrow Hom(\mathcal{F}, \mathcal{G}^*) \rightarrow (\mathcal{F}^*)^a \rightarrow (\mathcal{F}^*)^b.$$

Taking the dual we get

$$\mathcal{F}^a \rightarrow \mathcal{F}^b \rightarrow \text{Hom}(\mathcal{F}, \mathcal{G}^*)^* \rightarrow 0.$$

This time the term $\text{Hom}(\mathcal{F}, \mathcal{G}^*)^*$ is by definition the tensor product $\mathcal{F} \otimes_{\mathcal{O}} \mathcal{G}$.

In general there is a natural morphism

$$U \otimes_{\mathcal{O}} V = \text{Hom}(U, V^*)^* \rightarrow \text{Hom}(U^*, V) = U \otimes^{\mathcal{O}} V.$$

To define this map we define a trilinear morphism

$$\text{Hom}(U, V^*)^* \times U^* \times V^* \rightarrow \mathcal{O},$$

by $(\lambda, \mu, \nu) \mapsto \lambda(\nu\mu)$, the product $\nu\mu$ being the composed morphism

$$U \xrightarrow{\mu} \mathcal{O} \xrightarrow{\nu} V^*.$$

This trilinear map gives a morphism

$$\text{Hom}(U, V^*)^* \rightarrow \text{Hom}(U^*, (V^*)^*)$$

then note that $(V^*)^* = V$.

The above presentations for $\mathcal{F} \otimes^{\mathcal{O}} \mathcal{G}$ and $\mathcal{F} \otimes_{\mathcal{O}} \mathcal{G}$ (which are the same) are compatible with this natural morphism, so the natural morphism is an isomorphism

$$\mathcal{F} \otimes^{\mathcal{O}} \mathcal{G} \cong \mathcal{F} \otimes_{\mathcal{O}} \mathcal{G}.$$

□

We now come to the statement of the “relative Breen calculations in characteristic 0”. The case $S = \text{Spec}(k)$ for k a field is due to [9], and the relative case for a group scheme was suggested in [8] (where the case of cohomology with coefficients in the multiplicative group scheme was treated).

Theorem 9.2 *Suppose S is a scheme over $\text{Spec}(\mathbf{Q})$. Suppose V is a vector scheme over X and suppose $\mathcal{F}$ is a coherent sheaf over S . Then for n odd we have*

$$H^i(K(V/S, n)/S, \mathcal{F}) = \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{i/n} (V^*).$$

For n even we have

$$H^i(K(V/S, n)/S, \mathcal{F}) = \mathcal{F} \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}^{i/n} (V^*).$$

In both cases the answer is 0 if i/n is not an integer. The multiplicative structures on the left sides, in the case $\mathcal{F} = \mathcal{O}$, coincide with the obvious ones on the right sides. In the case of arbitrary $\mathcal{F}$, the natural structures of modules over the cohomology with coefficients in $\mathcal{O}$, on both sides, coincide.

We first recall the following result.

Proposition 9.3 *Suppose S is a scheme and V is a vector sheaf on S . Then the complexes*

$$\dots \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^j \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}^k V \rightarrow \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{j-1} \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}^{k+1} V \dots$$

and

$$\dots \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^j \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}^k V \rightarrow \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{j+1} \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}^{k-1} V \dots$$

are exact as sequences of vector sheaves (i.e. as sequences of sheaves on the site $Sch/\mathbf{C}$).

Proof: For $\mathcal{F} = \mathcal{O}$ this is Proposition 3.8 of [39]. The proof is easy, obtained by taking the graded symmetric powers of the cohomologically trivial complex $V \rightarrow V$ (placing this complex starting in odd or even degrees, leads to the two cases of the statement).

For a general $\mathcal{F}$ note that the cotensor product with a coherent sheaf is exact—indeed, for any vector sheaf U , $\text{Hom}(\mathcal{F}^*, U)$ is exact in U because $\mathcal{F}^*$ is represented by a scheme, i.e. an element of the site $Sch/\mathbf{C}$. The tensor product is equal to the cotensor product because both sides are coherent sheaves (here is where we use the hypothesis that V is a vector scheme, i.e. V^* is a coherent sheaf). $\square$

Proof of Theorem 9.2

Now we start the proof of Theorem 9.2. Suppose that it is true for $n \leq m-1$, and we prove it for $n = m$. (The case $m = 1$ to start the induction will be treated separately at the end.) Look at the fiber sequence

$$K(V/S, m-1) \rightarrow * \xrightarrow{p} K(V/S, m).$$

We will look at the Leray spectral sequence for the morphism p , for cohomology with coefficients in $\mathcal{F}$.

We may assume that S is affine

Let A_m denote the algebra $H^*(K(V/S, m)/S, \mathcal{O})$ (this is a sheaf of algebras over S) and let $A_m(\mathcal{F})$ denote the A_m -module $H^*(K(V/S, m)/S, \mathcal{F})$. Note that if $\mathcal{F}$ itself is an algebra-object then $A_m(\mathcal{F})$ is an A_m -algebra (graded-commutative). By the inductive hypothesis, the A_k and $A_k(\mathcal{F})$ are direct sums of coherent sheaves (coherent in each degree) for $k \leq m-1$.

The E_2 term of our spectral sequence is

$$H^i(K(V/S, m)/S, H^j(K(V/S, m-1)/S, \mathcal{F})) \Rightarrow H^{i+j}(*, \mathcal{F}).$$

In the case $\mathcal{F} = \mathcal{O}$ the E_2 term has a structure of algebra, and for arbitrary $\mathcal{F}$, a structure of module over that algebra. These are respectively

$$A_m(A_{m-1})$$

and

$$A_m(A_{m-1}(\mathcal{F})).$$

By induction we know that

$$A_{m-1}(\mathcal{F}) = A_{m-1} \otimes_{\mathcal{O}} \mathcal{F}.$$

The first possible nonzero differential in the spectral sequence is

$$H^i(K(V/S, m)/S, H^j(K(V/S, m-1)/S, \mathcal{F})) \rightarrow$$

$$H^{i+m}(K(V/S, m)/S, H^{j+1-m}(K(V/S, m-1)/S, \mathcal{F})).$$

We prove by a second induction on k , that for all $\mathcal{F}$ the answer is as given in the theorem, for $H^i(K(V/S, m)/S, \mathcal{F})$ for all $i \leq k$. Suppose this is true for $i \leq k-1$. Then the elements of the diagonal complex for the above differential, ending at $(i, j) = (k, 0)$, are all in the region $i \leq k-1$, except for the term $(k, 0)$. By our second inductive hypothesis (applied to the cohomology of $K(V/S, m)$ with coefficients in the coherent sheaves $H^j(K(V/S, m-1), \mathcal{F})$, this complex coincides with one of the two complexes appearing in Proposition 9.3, except maybe for the last term. However, the complex must be exact at the last stage because otherwise, what is left over would persist into E_∞ contradicting the answer of the spectral sequence (which must be $\mathcal{F}$ in degree 0 and 0 otherwise). Proposition 9.3 gives exactness of the complexes appearing there. Therefore the $E_2^{k,0}$ -term of our spectral sequence must also coincide with the last term of the complex from Proposition 9.3. For example in the case where m is odd, the end of the spectral sequence is

$$\begin{aligned} \dots &\rightarrow \mathcal{F} \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}^2(V^*) \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{(k-2m)/m} (V^*) \\ &\rightarrow \mathcal{F} \otimes_{\mathcal{O}} V^* \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{(k-m)/m} (V^*) \\ &\xrightarrow{d} E_2^{k,0}(\mathcal{F}) \rightarrow 0 \end{aligned}$$

(where we denote by d the last differential), whereas the end of the complex of Proposition 9.3 is

$$\dots \rightarrow \mathcal{F} \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}^2(V^*) \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{(k-2m)/m} (V^*)$$

$$\begin{aligned}
& \rightarrow \mathcal{F} \otimes_{\mathcal{O}} V^* \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{(k-m)/m} (V^*) \\
& \rightarrow \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{k/m} (V^*) \rightarrow 0.
\end{aligned}$$

Therefore if m is odd,

$$E_2^{k,0} = \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{k/n} (V^*).$$

The same holds with symmetric power instead of exterior power if m is even.

Cup product gives a bilinear morphism

$$\mu : E_2^{k-m,0} \times H^m(K(V/S, m), \mathcal{O}) = E_2^{k-m,0} \times V^* \rightarrow E_2^{k,0}.$$

Let

$$d' : \mathcal{F} \otimes_{\mathcal{O}} V^* \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{(k-2m)/m} (V^*) \rightarrow E_2^{k-m,0}(\mathcal{F})$$

denote the previous differential. We know by induction that d' establishes an isomorphism between $E_2^{k-m,0}(\mathcal{F})$ and

$$\mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{k/m} (V^*)$$

where this latter is considered as a quotient of the range of d' via Proposition 9.3.

Referring to the cup-product morphism μ considered above and its precursor

$$\mu' : E_2^{k-2m,m-1} \times V^* \rightarrow E_2^{k-m,m-1},$$

we have the Leibniz formula

$$d\mu'(a, v) = \mu(d'(a), v)$$

noting that the term V^* appearing in the formulas is $E_2^{m,0}$ so the differential acts trivially on the variable v .

We have exactly the same formula when the terms $E_2^{i,j}$ are replaced by their counterparts from the sequences of Proposition 9.3. Denote the multiplication in these counterparts by ν and ν' and the differentials by δ and δ' .

Call the isomorphism established by d ,

$$\psi^{k,0} : \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^{k/m} (V^*) \cong E_2^{k,0}$$

and similarly we have isomorphisms (by the inductive hypothesis) $\psi^{k-m,0}$, $\psi^{k-m,m-1}$ and $\psi^{k-2m,m-1}$. The definition of $\psi^{k,0}$ is given by the equation

$$\psi^{k,0}(\delta(b)) = d\psi^{k-m,m-1}(b).$$

Similarly we have

$$\psi^{k-m,0}(\delta'(b)) = d'\psi^{k-2m,m-1}(b).$$

We get

$$\psi^{k,0}(\nu(\delta'(a), v)) = \psi^{k,0}(\delta\nu'(a, v)) = d\psi^{k-m,m-1}(\nu'(a, v)).$$

On the other hand, in this region we know by induction that the ψ are compatible with products. Therefore we get

$$\begin{aligned} d\psi^{k-m,m-1}(\nu'(a, v)) &= d\mu'(\psi^{k-2m,m-1}(a), v) \\ &= \mu(d'\psi^{k-2m,m-1}(a), v) \\ &= \mu(\psi^{k-m,0}(\delta'(a)), v). \end{aligned}$$

In all we have established the formula

$$\psi^{k,0}(\nu(b, v)) = \mu(\psi^{k-m,0}(b), v)$$

for any $b = \delta'(a)$. But $\delta'(a)$ is surjective. This establishes the compatibility of our isomorphism $\psi^{k,0}$ with products (given already the compatibility of $\psi^{k-m,0}$).

We note in the above proof that elements of the tensor products are always (locally on S) finite sums of tensors. This can be seen for example from the proof of Lemma 9.1. Thus to check compatibility with products, for example, it suffices to check it on elementary tensors as we have done above.

This completes the proof of the theorem, modulo the case $m = 1$ which we now treat.

We have the fiber sequence

$$V \rightarrow S \xrightarrow{p} K(V, 1).$$

The higher direct images vanish for coefficients in a coherent sheaf $\mathcal{F}$ so the Leray spectral sequence implies that

$$R^i p_*(\mathcal{F}) = 0, \quad i > 0$$

and $R^0 p_*(\mathcal{F})$ is a local system on $K(V, 1)$ which when restricted to S gives

$$\mathcal{F}[V] \cong \mathcal{F} \otimes_{\mathcal{O}} \text{Sym}_{\mathcal{O}}(V^*).$$

Now we use the complex given in Proposition 9.3, which is basically a de Rham complex in our situation:

$$0 \rightarrow \mathcal{O} \rightarrow \mathrm{Sym}_{\mathcal{O}}(V^*) \rightarrow \mathrm{Sym}_{\mathcal{O}}(V^*) \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^i(V^*) \rightarrow \dots$$

which can then be tensored by $\mathcal{F}$ and remains exact. We can define the translation action of V on all of the terms, and the exact sequence remains an exact sequence of sheaves with action of V . All of the terms except for the first one are acyclic by the previous result. Therefore the cohomology of $K(V, 1)$ with coefficients in $\mathcal{F}$ is equal to the cohomology of the complex

$$\mathcal{F} \rightarrow \dots \rightarrow \mathcal{F} \otimes_{\mathcal{O}} \bigwedge_{\mathcal{O}}^i(V^*) \rightarrow \dots$$

One can check that the differentials are actually zero, so the cohomology is as desired.

One should check that the cup-product is equal to the obvious product structure on the exterior-algebra side of the answer. Instead of doing this (which as such would seem to be a difficult task) we proceed as follows. The above construction is functorial (contravariantly) for morphisms $V \rightarrow V'$. It is easy to see that if one considers an injection $V \hookrightarrow \mathcal{O}^a$ (which exists locally on S by the definition of vector scheme) then the morphism of functoriality induces a surjection on cohomology, coming from the surjection $\mathcal{O}^a \rightarrow V^*$. Thus to establish a formula for cup-products in cohomology, it suffices to establish the formula for the case $V = \mathcal{O}^a$. In that case we can apply the Künneth formula, or more precisely remark that the same Künneth formula holds for the cohomology as for the exterior algebra, and that these formulas are compatible via the above isomorphism. The Künneth formulae are both compatible with cup-products. Thus we can reduce to the case $V = \mathcal{O}$, but here the cohomology is concentrated in degrees 0 and 1 so there are no cup-products to verify (excepting the product with a degree 0 class but this case is easy). $\square$

Corollary 9.4 *Suppose S is a scheme and $T \rightarrow S$ is a relatively 1-connected very presentable n -stack. Then for any vector sheaf V on S ,*

$$H^i(T/S, V)$$

is a vector sheaf.

Proof: We use systematically (without further mention) the fact that the category of vector sheaves is closed under kernels, cokernels and extensions cf [39] [41]. The case of $T = K(U/S, n)$ for U a vector scheme and V a coherent sheaf is given by Theorem

9.2. The case of coefficients in any vector sheaf V is obtained by taking a resolution of V by coherent sheaves and using the long exact sequence of cohomology. The case of $T = K(U/S, n)$ for any vector sheaf U is obtained by taking a resolution of U by vector schemes (divided into two short exact sequences which give rise to two fibration sequences) and then applying the Leray spectral sequence. Finally, any relatively 1-connected very presentable T has a Postnikov tower (relative to S) whose stages are $K(U/S, n)$. Repeated application of the Leray spectral sequence gives the result. $\square$

Corollary 9.5 *If S is a scheme and $T \rightarrow S$ and $T' \rightarrow S$ are relatively 1-connected very presentable n -stacks then $\text{Hom}(T/S, T'/S)$ is very presentable.*

For this one has to use the fact that $\text{Aut}(V)$ is a very presentable group sheaf when V is a vector sheaf, see [41]. $\square$

The following application was the original motivation for Breen's calculations of the cohomology of the Eilenberg-MacLane sheaves [8] [9]. From our version Theorem 9.2, we obtain the corresponding result in the relative case in characteristic zero. Similar corollaries were stated for example for cohomology with coefficients in the multiplicative group $\mathbf{G}_m$, in [8].

Corollary 9.6 ([39] Corollary 3.11) *Suppose U, V are vector sheaves over a scheme S . Let $\text{Ext}_{\text{gp}}^i(U, V)$ denote the Ext sheaves between U and V considered as sheaves of abelian groups on Sch/S , let $\text{Ext}_{\text{vs}}^i(U, V)$ denote the Ext sheaves between U and V considered as vector sheaves on S . Then the natural morphisms are isomorphisms*

$$\text{Ext}_{\text{vs}}^i(U, V) \xrightarrow{\cong} \text{Ext}_{\text{gp}}^i(U, V).$$

The Ext^i vanish for $i > 2$.

Proof: Let $K.(U, n)$ denote the simplicial presheaf

$$Y \mapsto K.(U(Y), n)$$

given by the standard simplicial Eilenberg-MacLane construction (i.e. Dold-Puppe applied to the complex with U placed in degree n). We don't take the associated stack (as doing this or not doesn't affect the morphisms to an m -stack). Let $\mathbf{Z}K.(U, n)$ denote the associated presheaf of simplicial free abelian groups. Finally let $N\mathbf{Z}K.(U, n)$ be the presheaf of normalized complexes (in the homology direction i.e. with differential of degree -1) of this simplicial abelian group. For each Y ,

$$N\mathbf{Z}K.(U, n)(Y)$$

is a complex with homology group $U(Y)$ in degree n , and with all other cohomology groups equal to 0 in degrees $< 2n$. Thus if $\mathcal{F}$ is an injective sheaf of groups then the (cohomological) complex of sheaves

$$Hom(N\mathbf{Z}K.(U, n), \mathcal{F})$$

has homology sheaf $Hom(U, \mathcal{F})$ in degree n and zero homology in all other degrees $< 2n$. It follows that if $\mathcal{F}$ is any sheaf of groups then for $i < n$,

$$H^{n+i}Hom(N\mathbf{Z}K.(U, n), \mathcal{F}) = Ext_{\text{gp}}^i(U, \mathcal{F}).$$

On the other hand, this complex of sheaves also calculates $H^*(K(U, n), \mathcal{F})$. Thus we find that

$$H^{n+i}(K(U, n), \mathcal{F}) = Ext_{\text{gp}}^i(U, \mathcal{F}), \quad i < n.$$

This holds true for any sheaves of groups U and $\mathcal{F}$. This is one of the motivating observations of Breen's paper [9]—we have repeated the proof here for the reader's convenience.

Now suppose that U and $\mathcal{F}$ are vector sheaves. If U is a vector scheme and $\mathcal{F}$ is a coherent sheaf then, via the above observation, the relative Breen calculations (Theorem 9.2) show that $Ext_{\text{gp}}^i(U, \mathcal{F}) = 0$ for $i > 0$.

A coherent sheaf $\mathcal{F}$ is an injective object in the category of vector sheaves, and the functor $Hom(\cdot, \mathcal{F})$ is exact (cf the discussion of vector sheaves in [41] for example). Thus if U is any vector sheaf, we can (locally on S) resolve it by vector schemes and apply the previous paragraph. The functor $Ext^0(\cdot, \mathcal{F}) = Hom(\cdot, \mathcal{F})$ is exact (recall from [39] Lemma 3.2 or [41] Lemma 4.5 that morphisms of sheaves of abelian groups are the same as morphisms of vector sheaves so the Hom is the same in the two categories). Using this exactness we get that $Ext_{\text{gp}}^i(U, \mathcal{F}) = 0$ for $i > 0$. Finally, if V is any vector sheaf then we can resolve it by coherent sheaves, which is an injective resolution in the category of vector sheaves. This is also an acyclic resolution for Ext in the category of sheaves of abelian groups, so we obtain the isomorphism

$$Ext_{\text{vs}}^i(U, V) \xrightarrow{\cong} Ext_{\text{gp}}^i(U, V).$$

The vanishing of the Ext^i for $i > 2$ comes from the fact that any vector sheaf V has a resolution of length 2 (i.e. with terms in degrees 0, 1, 2) by coherent sheaves (cf [39] [41]). $\square$

We can apply 9.6 to the example discussed at the start of the appendix. Suppose S is a scheme and suppose $E^\cdot$ is a complex of vector bundles on S . The cohomology sheaves $V^i = \mathcal{H}^i(E^\cdot)$ are vector sheaves. In general, a complex with given cohomology objects is determined by higher extension classes in Ext^i for all values of $i \geq 2$. However, by

virtue of the above theorem the $Ext^i(V^j, V^k)$ vanish for $i \geq 3$. Thus the complex $E^\cdot$ is determined completely by the successive extension classes

$$\delta_{j,j+1} \in Ext^2(V^{j+1}, V^j).$$

The same is true for any complex of vector sheaves with V^j as cohomology objects.

Problem: describe the conditions which must be satisfied by the classes $\delta_{j,j+1}$ for the complex determined by these classes to be (quasiisomorphic to) a complex of vector bundles.

10. APPENDIX II: Representability of very presentable shape

The following result was stated without proof in ([39], the discussion above Theorem 5.7). Since we refer anew to this result in our discussion after Theorem 7.1 of the present paper, I felt it to be an opportune time to give a proof.

Theorem 10.1 *Suppose $\mathcal{F}$ is a connected n -stack on $Sch/\mathbf{C}$ such that the cohomology sheaves $H^i(\mathcal{F}, \mathcal{O})$ are represented by finite dimensional vector spaces. Suppose furthermore that $H^0(\mathcal{F}, \mathcal{O}) = \mathcal{O}$ and $H^1(\mathcal{F}, \mathcal{O}) = 0$. Then the $n+1$ -functor*

$$T \mapsto Hom(\mathcal{F}, T)$$

from 1-connected very presentable n -stacks of groupoids T to the same, is representable by a morphism $\mathcal{F} \rightarrow \Sigma$, with Σ being a 1-connected very presentable n -stack.

Proof: It suffices to have a morphism $\mathcal{F} \rightarrow \Sigma$ which induces an isomorphism

$$H^i(\Sigma, \mathcal{O}) \xrightarrow{\cong} H^i(\mathcal{F}, \mathcal{O})$$

for any i . We say that a morphism $\mathcal{F} \rightarrow \Sigma_m$ is m -arranged if the induced morphisms on cohomology with coefficients in $\mathcal{O}$ are isomorphisms for $k \leq i < m$ and injective for $k \leq i = m$. Note that the morphism $\mathcal{F} \rightarrow *$ is 1-arranged because of the hypothesis that $H^1(\mathcal{F}, \mathcal{O}) = 0$. The strategy of the proof (taken from E. Brown [11]) will be to suppose that we have constructed $\mathcal{F} \rightarrow \Sigma_m$ which is m -arranged. Then we will construct a factorization

$$\mathcal{F} \rightarrow \Sigma_{m+1} \rightarrow \Sigma_m$$

where the first morphism is $m+1$ -arranged. By induction this suffices to prove the theorem (we can stop as soon as we get to $m > n$).

So start with the situation of $\mathcal{F} \rightarrow \Sigma_m$, m -arranged, $m \geq 1$. Let

$$\mathcal{C} := Cone(\mathcal{F} \rightarrow \Sigma_m)$$

so we have a map $\Sigma_m \rightarrow \mathcal{C}$ which restricted to $\mathcal{F}$ gives a map homotopic to the basepoint $*$ $\rightarrow \mathcal{C}$ (this basepoint is included in the definition of *Cone*—it is the vertex of the cone over $\mathcal{F}$).

It is easy to see using the m -arrangedness of our map, that the cohomology of $\mathcal{C}$ with coefficients in $\mathcal{O}$ vanishes in degrees $\leq m$. Furthermore, the $m+1$ -st cohomology fits into a long exact sequence with those of $\mathcal{F}$ and Σ_m :

$$\begin{aligned} 0 &\rightarrow H^m(\Sigma_m, \mathcal{O}) \rightarrow H^m(\mathcal{F}, \mathcal{O}) \\ &\rightarrow H^{m+1}(\mathcal{C}, \mathcal{O}) \rightarrow H^{m+1}(\Sigma_m, \mathcal{O}) \\ &\rightarrow H^{m+1}(\mathcal{F}, \mathcal{O}) \rightarrow H^{m+2}(\mathcal{C}, \mathcal{O}) \\ &\rightarrow H^{m+2}(\Sigma_m, \mathcal{O}) \rightarrow \dots \end{aligned}$$

The cohomology of Σ_m with coefficients in $\mathcal{O}$ is a finite dimensional vector space, by Theorem 9.2 (this case is contained in the original characteristic 0 version obtainable from [9]). The property of being represented by a finite dimensional vector space is closed under extensions, kernels and cokernels ([39] Theorem 3.3 and [41] Corollary 4.10 and Theorem 4.11). Therefore the cohomology of $\mathcal{C}$ with coefficients in $\mathcal{O}$ is again a (sheaf represented by a) finite dimensional vector space.

Now let

$$W^* := H^{m+1}(\mathcal{C}, \mathcal{O})$$

define the finite dimensional vector space W . We get a morphism $\mathcal{C} \rightarrow K(W, m+1)$ which is universal for morphisms to $K(U, m+1)$ with U a finite dimensional vector space. In particular it induces an isomorphism

$$H^{m+1}(K(W, m+1), \mathcal{O}) \xrightarrow{\cong} H^{m+1}(\mathcal{C}, \mathcal{O}).$$

We will compare the previous long exact sequence with the long exact sequence that occurs at the start of the Leray-Serre spectral sequence for the morphism $p : \Sigma_m \rightarrow K(W, m+1)$. Set

$$Fib := Fib(\Sigma_m \rightarrow K(W, m+1)).$$

Note that we have a morphism $\mathcal{F} \rightarrow Fib$.

The higher direct images occuring in the Leray-Serre spectral sequence for p are constant local systems over the base, because $K(W, m+1)$ is 1-connected. In other words,

$$R^i p_* \mathcal{O} = H^i(Fib, \mathcal{O}) \times K(W, m+1) \rightarrow K(W, m+1).$$

Do a standard type of spectral sequence argument. First of all, for $k < m$ we prove by induction on k that for all $i \leq k$, the $H^i(Fib, \mathcal{O})$ are finite dimensional vector spaces.

Suppose we know this for $k - 1$. Then in view of the vanishing of the cohomology of $K(W, m + 1)$ with vector space coefficients (the sheaves represented by vector spaces are $\mathcal{O}^a$) the terms $E_2^{i,j}$ with $1 \leq i \leq m$ and $j < k$ vanish; whereas for $j = 0$ we have

$$E_2^{i,0} = H^i(Fib, \mathcal{O})$$

because of the fact that $K(W, m + 1)$ is 1-connected. Therefore the terms $E_2^{k,0}$ persist to E_∞ and we have

$$H^k(Fib, \mathcal{O}) = H^k(\Sigma_m, \mathcal{O}) = H^i(\mathcal{F}, \mathcal{O}).$$

In view of the hypothesis of the theorem, this proves the induction step of the first part of the argument. Incidentally we get that the morphism $\mathcal{F} \rightarrow Fib$ induces an isomorphism on cohomology with coefficients in $\mathcal{O}$ in degrees $k < m$. Now look at the term $E_2^{m,0} = H^m(Fib, \mathcal{O})$. The only differential concerning it is

$$d_{m+1} : H^m(Fib, \mathcal{O}) \rightarrow H^{m+1}(K(W, m + 1), \mathcal{O})$$

(noting that we already have $H^0(Fib, \mathcal{O}) = \mathcal{O}$). From our hypothesis which implies that $H^1(Fib, \mathcal{O}) = 0$ we get, similarly, that the only differential concerning the term $E_2^{m+1,0}$ is

$$d_{m+2} : H^{m+1}(Fib, \mathcal{O}) \rightarrow H^{m+2}(K(W, m + 1), \mathcal{O}).$$

From these and the fact that the spectral sequence abuts to the cohomology of Σ_m , we get the long exact sequence

$$\begin{aligned} 0 &\rightarrow H^m(\Sigma_m, \mathcal{O}) \rightarrow H^m(Fib, \mathcal{O}) \\ &\rightarrow H^{m+1}(K(W, m + 1), \mathcal{O}) \rightarrow H^{m+1}(\Sigma_m, \mathcal{O}) \rightarrow \\ &H^{m+1}(Fib, \mathcal{O}) \rightarrow H^{m+2}(K(W, m + 1), \mathcal{O}) \rightarrow H^{m+2}(\Sigma_m, \mathcal{O}). \end{aligned}$$

Remark that $H^{m+2}(K(W, m + 1), \mathcal{O}) = 0$ —this comes from Theorem 9.2 and it is here where we use $m \geq 1$. In particular, the morphism

$$H^{m+2}(K(W, m + 1), \mathcal{O}) \rightarrow H^{m+2}(\mathcal{C}, \mathcal{O})$$

is injective for the trivial reason that the left side is 0. Recall that the same induced morphism in degree $m + 1$ was an isomorphism (by the construction of W). Therefore, comparing with the previous long exact sequence and using the 5-lemma, we get that the morphism

$$\mathcal{F} \rightarrow Fib$$

induces isomorphisms on cohomology in degrees $\leq m$ and an injection in degree $m + 1$. In other words this morphism is $m + 1$ -arranged. Thus we can set

$$\Sigma_{m+1} := Fib$$

and we have completed our inductive construction to prove the theorem. $\square$

Definition: If $\mathcal{F}$ satisfies the condition of Theorem 10.1 then we obtain the representing 1-connected very presentable $\Sigma(\mathcal{F})$ with universal morphism

$$\mathcal{F} \rightarrow \Sigma(\mathcal{F}).$$

We define (for any basepoint $f : Y \rightarrow \mathcal{F}$)

$$\pi_i^{\text{vp}}(\mathcal{F} \times Y/Y, f) := \pi_i(\Sigma \times Y/Y, f).$$

In the latter case we usually just take a basepoint $f \in \mathcal{F}(\text{Spec } \mathbf{C})$ and then denote this by $\pi_i^{\text{vp}}(\mathcal{F}, f)$.

Theorem 10.2 *Suppose $\mathcal{F}$ and $\mathcal{G}$ are n -stacks with basepoint $g \in \mathcal{G}(\text{Spec } \mathbf{C})$, which satisfy the criterion of Theorem 10.1 so their shapes are representable. Suppose*

$$f : \mathcal{F} \rightarrow \mathcal{G}$$

is a morphism of n -stacks with the following property (we denote by $\mathcal{H}$ the fiber over g): the local systems $R^i f_(\mathcal{O})$ are isomorphic to $\mathcal{O}^{a_i}$ on $\mathcal{G}$, and that the morphisms*

$$R^i f_*(\mathcal{O})|_g \rightarrow H^i(\mathcal{H}, \mathcal{O})$$

are isomorphisms. Suppose that

$$H^1(\mathcal{F}, \mathcal{O}) = H^1(\mathcal{G}, \mathcal{O}) = H^1(\mathcal{H}, \mathcal{O}) = \mathcal{O}$$

and

$$H^1(\mathcal{F}, \mathcal{O}) = H^1(\mathcal{G}, \mathcal{O}) = H^1(\mathcal{H}, \mathcal{O}) = 0.$$

Then we have a fiber sequence for the representing objects

$$\Sigma(\mathcal{H}) \rightarrow \Sigma(\mathcal{F}) \rightarrow \Sigma(\mathcal{G}).$$

Proof: The Leray spectral sequence for f is

$$H^i(\mathcal{G}, R^j f_*(\mathcal{O})) \Rightarrow H^{i+j}(\mathcal{F}, \mathcal{O}).$$

In view of the hypothesis, this becomes

$$H^i(\mathcal{G}, \mathcal{O}) \otimes_{\mathcal{O}} H^j(\mathcal{H}, \mathcal{O}) \Rightarrow H^{i+j}(\mathcal{F}, \mathcal{O}).$$

On the other hand, we obtain a morphism of representing shapes

$$\Sigma(\mathcal{F}) \rightarrow \Sigma(\mathcal{G})$$

(by the universal property of $\Sigma(\mathcal{F})$). Let Fib denote the fiber of $\Sigma(\mathcal{F})$ (over the image of the point g). Note that the $R^i\Sigma(f)_*(\mathcal{O})$ are constant on $\Sigma(\mathcal{G})$ because $\Sigma(\mathcal{G})$ is 1-connected. In particular

$$R^i f_*(\mathcal{O})|_g \xrightarrow{\cong} H^i(Fib, \mathcal{O}).$$

We obtain the spectral sequence

$$H^i(\Sigma(\mathcal{G}), \mathcal{O}) \otimes_{\mathcal{O}} H^j(Fib, \mathcal{O}) \Rightarrow H^{i+j}(\Sigma(\mathcal{F}), \mathcal{O}).$$

Note that the $\pi_i(Fib)$ are finite dimensional vector spaces (using the long exact sequence of homotopy groups of a fibration, and [41] Theorem 4.11 applied to the case of trivial base $S = *$). Thus $H^j(Fib, \mathcal{O})$ are finite dimensional vector spaces.

The composition

$$\mathcal{H} \rightarrow \Sigma(\mathcal{F}) \rightarrow \Sigma(\mathcal{G})$$

is homotopic to the constant map at the basepoint, so we get a map

$$\mathcal{H} \rightarrow Fib.$$

This gives maps $H^i(Fib, \mathcal{O}) \rightarrow H^i(\mathcal{H}, \mathcal{O})$. We claim that these are isomorphisms, which would imply that $\mathcal{H} \rightarrow Fib$ is a map representing the very presentable shape of $\mathcal{H}$, in other words $\Sigma(\mathcal{H}) \cong Fib$, thus giving the desired result.

To prove the claim, note that the maps in question are compatible via the previous identifications, with the maps

$$R^i\Sigma(\mathcal{F})_*\mathcal{O} \rightarrow R^i f_*(\mathcal{O}).$$

These in turn fit into a morphism of Leray spectral sequences. We show using a spectral sequence argument, by induction on k , that for all $i \leq k$ we have

$$H^i(Fib, \mathcal{O}) \xrightarrow{\cong} H^i(\mathcal{H}, \mathcal{O})$$

or equivalently

$$R^i\Sigma(\mathcal{F})_*\mathcal{O} \xrightarrow{\cong} R^i f_*(\mathcal{O}).$$

Suppose this is true for $k - 1$. Then look at the term $E_2^{0,k} = H^k(\mathcal{H}, \mathcal{O})$. When we look at the r th differential

$$d_r^{0,k} : E_r^{0,k} \rightarrow E_r^{r,k+1-r}$$

the term $E_r^{r,k+1-r}$ has not yet been touched by any term $E^{i,j}$ with $j \geq k$, and after this differential, the term $E^{r,k+1-r}$ is no longer touched by any further differentials. We have a morphism of spectral sequences (the above remarks apply to both) which induces an isomorphism on the abutments. It follows that the morphism between spectral sequences induces an isomorphism on images of $d_r^{0,k}$. Furthermore, the morphism induces an isomorphism on the intersection (for all r) of the kernels of the $d_r^{0,k}$. This implies that the morphism induces an isomorphism on $E_2^{0,k}$ and we obtain the inductive step for k . This proves the claim and hence the theorem. $\square$

Corollary 10.3 *Suppose Y is a simply connected finite CW complex. Let $\mathcal{F}$ be the constant n -stack associated to the constant prestack with values Y (or more precisely, with values the n -type $\tau_{\leq n}Y$). Fix a basepoint $y \in Y$ which also gives a basepoint section of $\mathcal{F}$. Let $\mathcal{F} \rightarrow \Sigma$ be the morphism representing the shape of $\mathcal{F}$. The morphisms induced by $Y \rightarrow \Sigma(\text{Spec } \mathbf{C})$,*

$$\pi_i(Y, y) \rightarrow \pi_i(\Sigma, y)$$

induce isomorphisms

$$\pi_i(Y, y) \otimes_{\mathbf{Z}} \mathcal{O} \cong \pi_i(\Sigma, y).$$

Proof: Using the previous theorem, we can reduce by the Postnikov tower to the case $Y = K(A, n)$ for a finitely generated abelian group A . Then the Breen calculations imply that the morphism

$$K(A, n) \rightarrow K(A \otimes_{\mathbf{Z}} \mathcal{O}, n)$$

induces an isomorphism on cohomology with coefficients in $\mathcal{O}$. This implies that

$$\Sigma(K(A, n)) = K(A \otimes_{\mathbf{Z}} \mathcal{O}, n),$$

which gives the statement of the corollary. $\square$

Definition: Fix n . If Y is a 1-connected finite CW complex, then we define the *complexification* of Y denoted $Y \otimes \mathbf{C}$ to be the n -stack $\Sigma(\mathcal{F})$ representing the very presentable shape of the constant n -stack $\mathcal{F}$ with values $\tau_{\leq n}Y$. Note that this notion depends on n because we have chosen not to treat the questions arising if we try to take $n = \infty$.

Example: If we apply this to $Y = S^2$ then we obtain $\Sigma = S^2 \otimes \mathbf{C}$ as defined in §6 above. This is easy to see because, using the previous theorem, the homotopy sheaves of Σ are $\mathcal{O}$ in degrees 2 and 3; then there are only two possibilities for Σ and they are distinguished by the vanishing or nonvanishing of the Whitehead product. As the Whitehead product is nonzero for S^2 and the isomorphisms of the previous theorem are compatible with the Whitehead product (exercise), this implies that the Whitehead product for Σ is nontrivial, therefore Σ is equal to the T defined in §6.

We propose the above results as a way of interpreting what it means to look at the “complexified homotopy type of a space Y ”. We could do the same thing over the ground field $\mathbf{Q}$, and then we propose that this is what it means to look at the “rational homotopy type” of Y . This notion is preserved by base extension of the ground field.

Of course this should all be related to the usual definitions of Quillen, Sullivan, Morgan, Hain et.al. which refer (excepting Quillen) to algebras of differential forms. In those theories, base extension is obtained by tensoring the algebra of forms with the field extension. It has always been somewhat unclear what geometric interpretation to put on this base-extension process, and we propose the above theory as a way of obtaining a reasonable interpretation. We don’t, however, get into details of the relationship between the above theory and the differential-forms theories.

One advantage of the present formulation is that it explains what is going on in the non-simply connected case: the shape of the constant sheaf $\mathcal{F} = \text{Const}(Y)$ is no longer representable by a very presentable object (except in fairly restricted cases such as finite fundamental group). Thus, the object which carries the “rational homotopy” information of Y is the shape itself, rather than the representing object which may not exist. The shape, i.e. the functor

$$T \mapsto \text{Hom}(\mathcal{F}, T)$$

exists even when Y is not simply connected.

Proof of Theorem 2.1

The statement of Theorem 2.1 from §2 is very slightly different from the statement 10.1 given above. We indicate here how to get 2.1. Suppose that $\mathcal{F}$ is an n -stack on $\text{Sch}/\mathbf{C}$ such that for any affine algebraic group G ,

$$K(G, 1) \xrightarrow{\cong} \text{Hom}(\mathcal{F}, K(G, 1)).$$

In particular this implies that $H^0(\mathcal{F}, \mathcal{O}) = \mathcal{O}$ and $H^1(\mathcal{F}, \mathcal{O}) = 0$. With the hypothesis that $H^i(\mathcal{F}, \mathcal{O})$ are represented by finite dimensional vector spaces, we can apply Theorem 10.1 to get a morphism

$$\mathcal{F} \rightarrow \Sigma(\mathcal{F})$$

universal for morphisms to 1-connected very prepresentable T . Note that $\Sigma(\mathcal{F})$ is 1-connected. We have to show that it is also universal for morphisms to 0-connected very presentable T ; suppose that T is one such. We may choose a basepoint t , and let $G = \pi_1(T, t)$ (which is an affine algebraic group). We have a fiber sequence

$$T' \rightarrow T \rightarrow K(G, 1).$$

This gives a diagram

$$\begin{array}{ccccc} \mathrm{Hom}(\Sigma(\mathcal{F}), T') & \rightarrow & \mathrm{Hom}(\Sigma(\mathcal{F}), T) & \rightarrow & \mathrm{Hom}(\Sigma(\mathcal{F}), K(G, 1)) \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{Hom}(\mathcal{F}, T') & \rightarrow & \mathrm{Hom}(\mathcal{F}, T) & \rightarrow & \mathrm{Hom}(\mathcal{F}, K(G, 1)), \end{array}$$

where the horizontal sequences are fiber sequences. Since $\Sigma(\mathcal{F})$ is 1-connected we have

$$K(G, 1) \xrightarrow{\cong} \mathrm{Hom}(\Sigma(\mathcal{F}), K(G, 1)),$$

and the same holds for $\mathcal{F}$ by hypothesis. Therefore the vertical map on the right is an equivalence between $K(G, 1)$ and our diagram becomes

$$\begin{array}{ccccc} \mathrm{Hom}(\Sigma(\mathcal{F}), T') & \rightarrow & \mathrm{Hom}(\Sigma(\mathcal{F}), T) & \rightarrow & K(G, 1) \\ \downarrow & & \downarrow & & \downarrow = \\ \mathrm{Hom}(\mathcal{F}, T') & \rightarrow & \mathrm{Hom}(\mathcal{F}, T) & \rightarrow & K(G, 1). \end{array}$$

Now note that T' is a 1-connected very presentable n -stack, so the vertical arrow on the left is an equivalence. Since the base $K(G, 1)$ is 0-connected, we can use the long exact sequences of homotopy for these fibrations to conclude that the vertical morphism in the middle is an equivalence. This is what we needed to know to establish the universal property of $\mathcal{F} \rightarrow \Sigma(\mathcal{F})$ for Theorem 2.1. $\square$

A relative version

While we are on the subject, we give a relative version of Theorem 10.1. Recall [39] [41] that if Y is a scheme then a 1-connected n -stack $\mathcal{F} \rightarrow Y$ (which can also be thought of as a 1-connected n -stack on the site Sch/Y of schemes over Y) is said to be *very presentable* if for any basepoint section $f : Y' \rightarrow \mathcal{F}$ for a scheme $Y' \rightarrow Y$, the homotopy group sheaves $\pi_i(\mathcal{F}|_{Y'}, f)$ are vector sheaves on Y' . Since (for the present discussion) we have assumed $\mathcal{F}$ to be relatively 1-connected, the homotopy group sheaves don't depend on the choice of basepoint (indeed, the choice of basepoint is locally unique up to homotopy which itself is unique up to—nonunique—homotopy). Therefore they descend to sheaves of abelian groups $\pi_i(\mathcal{F}/Y)$ on Y . For $\mathcal{F}$ to be very presentable, it is equivalent to require that these be vector sheaves.

We introduce the following terminology. We say that a covariant endofunctor F from the category of vector sheaves on Y to itself, is *anchored* if the natural map

$$F(U) \rightarrow \mathrm{Hom}(\mathrm{Hom}(F(\mathcal{O}), \mathcal{O}), U)$$

is an isomorphism for any coherent sheaf U (recall that the coherent sheaves are the injective objects in the category of vector sheaves). The above natural map comes from the trilinear map

$$F(U) \times \text{Hom}(F(\mathcal{O}), \mathcal{O}) \times \text{Hom}(U, \mathcal{O}) \rightarrow \mathcal{O}$$

defined by $(f, g, h) \mapsto g(F(h)(f))$.

Lemma 10.4 (A) *If*

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$$

is a short exact sequence of natural transformations between covariant endofunctors on the category of vector sheaves over Y , then if any two of the three endofunctors is anchored, so is the third.

(B) *If F is an anchored endofunctor which is also left exact, then F is representable $F(V) = \text{Hom}(W, V)$ for a vector sheaf $W = \text{Hom}(F(\mathcal{O}), \mathcal{O})$.*

Proof: (A) follows from the 5-lemma. For (B) suppose F is a left exact anchored endofunctor. Set $W := \text{Hom}(F(\mathcal{O}), \mathcal{O})$. The natural map $F(U) \rightarrow \text{Hom}(W, U)$ is an isomorphism for coherent sheaves U . On the other hand, both sides are left exact in U . Suppose

$$0 \rightarrow U \rightarrow U' \rightarrow U''$$

is an exact sequence with U' and U'' being coherent sheaves. Then we obtain exact sequences

$$0 \rightarrow F(U) \rightarrow F(U') \rightarrow F(U'')$$

and

$$0 \rightarrow \text{Hom}(W, U) \rightarrow \text{Hom}(W, U') \rightarrow \text{Hom}(W, U''),$$

and our natural map is a morphism between these exact sequences inducing isomorphisms on the last two terms. Thus $F(U) \rightarrow \text{Hom}(W, U)$ is an isomorphism. This completes the proof in view of the fact that (locally on S) any vector sheaf U fits into such a short exact sequence. $\square$

Lemma 10.5 *Suppose V is a vector sheaf. Then the endofunctor on the category of vector sheaves defined by*

$$U \mapsto H^i(K(V, m), U)$$

is anchored.

Proof: This follows immediately from Theorem 9.2 if V is a vector scheme. Now suppose that we have an exact sequence

$$0 \rightarrow V' \rightarrow V'' \rightarrow V \rightarrow 0$$

where V'' is a vector scheme, and where we know the lemma for V' . This gives a fibration sequence

$$K(V'', m) \rightarrow K(V, m) \rightarrow K(V', m+1),$$

and taking the cohomology with coefficients in a coherent sheaf U leads to a Leray spectral sequence

$$H^i(K(V', m+1), H^j(K(V'', m), U)) \Rightarrow H^{i+j}(K(V, m), U).$$

The cohomology of the fiber are again coherent sheaves by Theorem 9.2, so by the lemma for V' the natural map occurring in the definition of “anchored” induces an isomorphism on the E_2 terms of the spectral sequence. Since the property of being anchored is preserved by kernels, cokernels and extensions, we get that the cohomology of $K(V, m)$ is anchored. $\square$

Corollary 10.6 *Suppose T is a relatively 1-connected very presentable n -stack over a scheme S . Then the endofunctor*

$$U \mapsto H^i(T/S, U)$$

is anchored.

Proof: Decompose T into a Postnikov tower where the pieces are of the form $K(V, m)$ for vector sheaves V $\square$

Theorem 10.7 *Suppose S is a scheme and $\mathcal{F} \rightarrow S$ is a morphism of n -stacks on $Sch/\mathbf{C}$. Suppose that for any vector sheaf V over S , the cohomology $H^i(\mathcal{F}/S, V)$ is again a vector sheaf over S . Suppose furthermore that $H^0(\mathcal{F}/S, V) = V$ and $H^1(\mathcal{F}/S, V) = 0$ for any vector sheaf V . Finally suppose that the functors $V \mapsto H^i(\mathcal{F}/S, V)$ are anchored. Then the functor*

$$T \mapsto Hom(\mathcal{F}/S, T/S)$$

from relatively 1-connected very presentable n -stacks of groupoids $T \rightarrow S$ to the same, is represented by a morphism $\mathcal{F} \rightarrow \Sigma$ over S , with $\Sigma \rightarrow S$ being a relatively 1-connected and very presentable n -stack over S .

Proof: Follow the same outline as for the proof of Theorem 10.1. We try to find a relatively 1-connected very presentable $\Sigma \rightarrow Y$ with a morphism

$$\mathcal{F} \rightarrow \Sigma$$

inducing an isomorphism on cohomology with coefficients in any coherent sheaf U on S (the isomorphism for coefficients in any vector sheaf U then follows by resolving U by coherent sheaves).

We say that a morphism $\mathcal{F} \rightarrow \Sigma_m$ is *m-arranged* if the induced morphisms on cohomology with coefficients in any coherent sheaf U on S are isomorphisms for $k \leq i < m$ and injective for $k \leq i = m$. Note that the morphism $\mathcal{F} \rightarrow S$ is 1-arranged because of the hypothesis that $H^1(\mathcal{F}/S, U) = 0$. Thus we may take $\Sigma_1 := S$. The strategy of the proof will be to suppose for some $m \geq 1$ that we have constructed $\mathcal{F} \rightarrow \Sigma_m$ which is *m-arranged*. Then we will construct a factorization

$$\mathcal{F} \rightarrow \Sigma_{m+1} \rightarrow \Sigma_m$$

where the first morphism is $m + 1$ -arranged. By induction this suffices to prove the theorem.

Make the same constructions, using the same notations (which we won't repeat here) as in the proof of Theorem 10.1. Along the way, replace the cohomology with coefficients in $\mathcal{O}$ (and the higher direct images of $\mathcal{O}$ etc.) by cohomology with coefficients in any coherent sheaf U on S .

We obtain the first long exact sequence (actually valid for any vector sheaf U as coefficients)

$$\begin{aligned} 0 &\rightarrow H^m(\Sigma_m/S, U) \rightarrow H^m(\mathcal{F}/S, U) \\ &\rightarrow H^{m+1}(\mathcal{C}/S, U) \rightarrow H^{m+1}(\Sigma_m/S, U) \\ &\rightarrow H^{m+1}(\mathcal{F}/S, U) \rightarrow H^{m+2}(\mathcal{C}/S, U) \\ &\rightarrow H^{m+2}(\Sigma_m/S, U) \rightarrow \dots \end{aligned}$$

The cohomology of Σ_m with coefficients in a vector sheaf is again a vector sheaf, by Corollary 9.4 above. The property of being represented by a finite dimensional vector space is closed under extensions, kernels and cokernels ([39] Theorem 3.3 and [41] Corollary 4.10 and Theorem 4.11). Therefore the cohomology of $\mathcal{C}$ with coefficients in a vector sheaf U is again a vector sheaf.

When we come to the construction of W we need to say something more—this is the reason for introducing the notion of “anchored” above. The functor

$$U \mapsto H^{m+1}(\mathcal{C}, U)$$

is anchored. This comes from the facts that the cohomology of Σ_m is anchored by Corollary 10.6, that the cohomology of $\mathcal{F}$ is anchored by hypothesis, and the fact that being anchored is preserved by kernels, cokernels and extensions (Lemma 10.4). On the other hand, the fact that the cohomology of $\mathcal{C}$ vanishes in degrees $0 < i \leq m$ (note that $m \geq 1$) implies that the above functor is left-exact. Therefore by Lemma 10.4 (B) it is representable by a vector sheaf W : we have

$$H^{m+1}(\mathcal{C}, U) = \text{Hom}(W, U).$$

In particular there is a tautological class in $H^{m+1}(\mathcal{C}, W)$ corresponding to a morphism $\mathcal{C} \rightarrow K(W, m+1)$, and this morphism is universal for morphisms from $\mathcal{C}$ to things of the form $K(U, m+1)$. In particular it induces an isomorphism

$$H^{m+1}(K(W, m+1), U) \xrightarrow{\cong} H^{m+1}(\mathcal{C}, U).$$

Again set

$$\text{Fib} := \text{Fib}(\Sigma_m \rightarrow K(W, m+1)).$$

Note that we have a morphism $\mathcal{F} \rightarrow \text{Fib}$.

Compare the first long exact sequence with the long exact sequence that occurs at the start of the Leray-Serre spectral sequence for the morphism $p : \Sigma_m \rightarrow K(W, m+1)$, using the same argument as in the proof of Theorem 10.1. We need to know that the morphism

$$H^{m+2}(K(W, m+1), U) \rightarrow H^{m+2}(\mathcal{C}, U)$$

is injective for cohomology with coefficients in a coherent sheaf U (recall that only coherent sheaves occur as coefficients for the cohomology in the definition of arrangedness—one goes back to the general case after the induction on m is finished). To prove this we again show that $H^{m+2}(K(W, m+1), U) = 0$ (note that this wouldn't be true if U were not a coherent sheaf and that is the reason why we restrict to coherent sheaves in the definition of arrangedness). In fact, using that $m \geq 1$ and following the argument of 10.1 we get that

$$H^{m+2}(K(W, m+1), U) = \text{Ext}^1(W, U).$$

However, a coherent sheaf U is an injective object in the category of vector sheaves [41] Lemma 4.17, so $\text{Ext}^1(W, U) = 0$. This gives a proof of the desired statement. Alternatively one can obtain a proof using a spectral sequence argument with a resolution

$$0 \rightarrow V \rightarrow V' \rightarrow V'' \rightarrow W \rightarrow 0$$

of W by vector schemes (decompose this into two short exact sequences and use a Leray spectral sequence argument for each of the corresponding fibration sequences).

After that the rest of the argument works exactly the same way as in Theorem 10.1 (calling upon Theorem 9.2 in the relative case as necessary). We don't repeat this. $\square$

References

- [1] J. Adams. *Infinite Loop Spaces*, Princeton University Press *Annals of Math. Studies* **90** (1978).
- [2] M. Artin. Versal deformations and algebraic stacks, *Inventiones Math.* **27** (1974), 165-189.
- [3] J. Baez, J. Dolan. Higher dimensional algebra and topological quantum field theory. Preprint q-alg (95-03).
- [4] J. Bénabou. *Introduction to Bicategories*, Lect. Notes in Math. **47**, Springer-Verlag (1967).
- [5] P. Berthelot. Cohomologie cristalline des schémas de caractéristique $p > 0$. Springer *Lecture Notes in Math.* **407** (1974).
- [6] I. Biswas. Secondary invariants of Higgs bundles. *International J. of Math.* **6** (1995), 193-204.
- [7] S. Bloch, H. Esnault. Algebraic Chern-Simons theory. Preprint alg-geom 9602001.
- [8] L. Breen. Extensions of abelian sheaves and Eilenberg-MacLane algebras. *Invent. Math.* **9** (1969), 15-44.
- [9] L. Breen. Extensions du groupe additif. *Publ. Math. I.H.E.S.* **48** (1978), 39-126.
- [10] L. Breen. On the classification of 2-gerbs and 2-stacks. *Astérisque* **225**, Soc. Math. de France (1994).
- [11] E. Brown, Jr. Finite computability of Postnikov complexes. *Ann. of Math.* **65** (1957), 1-20.
- [12] K. Brown. Abstract homotopy theory and generalized sheaf cohomology. *Trans. A.M.S.* **186** (1973), 419-458.
- [13] K. Brown, S. Gersten. *Algebraic K-theory as generalize sheaf cohomology* Springer *Lecture Notes in Math.* **341** (1973), 266-292.
- [14] J. Carlson, H. Clemens, J. Morgan. On the mixed Hodge structure associated to π_3 of a simply connected complex projective manifold. *Annales E.N.S.* **14** (1981) 323-338.

- [15] J.-M. Cordier, T. Porter. Homotopy-coherent category theory. *Trans. Amer. Math. Soc.* **349** (1997), 1-54.
- [16] P. Deligne, D. Mumford. The irreducibility of the space of curves of a given genus, *Publ. Math. I.H.E.S.* **36**, (1969), 75-110.
- [17] H. Esnault, E. Viehweg. Higher Kodaira-Spencer classes. Preprint.
- [18] J. Giraud. *Cohomologie nonabelienne* Grundlehren **179** Springer-Verlag (1971).
- [19] R. Gordon, A.J. Power, R. Street. Coherence for tricategories *Memoirs A.M.S.* **117** (1995), 558 ff.
- [20] A. Grothendieck. *Pursuing Stacks*, unpublished manuscript.
- [21] A. Hirschowitz. Cohérence et dualité sur le gros site de Zariski, *Algebraic curves and projective geometry* (Ballico, Ciliberto eds.), *Lecture Notes in Math.* **1389**, (1989), 91-102.
- [22] N. Hitchin. Stable bundles and integrable systems. *Duke Math. J.* **54** (1987), 91-114.
- [23] N. Hitchin. The self-duality equations on a Riemann surface. *Proc. London Math. Soc.* **55** (1987), 59-126.
- [24] L. Illusie. *Le complexe cotangent I et II*, *Lecture Notes in Math.* **239**, **283**, Springer-Verlag, Berlin (1971,1972).
- [25] Infinitesimal variation of Hodge structure, *Compositio* **50** (1983):
IVHS (I): J. Carlson, M. Green, P. Griffiths, J. Harris, 109-205;
IVHS (II): P. Griffiths, J. Harris: Infinitesimal invariant of Hodge classes, 207-265;
IVHS (III): P. Griffiths: Determinantal varieties and the infinitesimal invariant of normal functions, 267-324.
- [26] D. Jaffe. Coherent functors and torsion in the Picard group. *Trans. Amer. Math. Soc.* **349** (1997), 481-527.
- [27] J.F. Jardine. Simplicial presheaves, *J. Pure and Appl. Algebra* **47** (1987), 35-87.
- [28] A. Joyal, Letter to A. Grothendieck (referred to in Jardine's paper).
- [29] Y. Karpishpan. Higher-order differentials of the period map and higher Kodaira-Spencer classes. Preprint alg-geom 9405005.

- [30] Y. Karpishpan. Infinite-dimensional Lie algebras and the period map for curves. Preprint alg-geom 9406006.
- [31] G. Laumon, L. Moret-Bailly. Champs algébriques. Preprint, Orsay **42** (1992).
- [32] O. Leroy. Sur une notion de 3-catégorie adaptée à l'homotopie. Preprint Univ. de Montpellier 2 (1994).
- [33] D. Quillen. Rational homotopy theory. *Ann. of Math.* **90** (1969), 205-295.
- [34] Z. Ran. Derivatives of moduli. *Int. Math. Res. Notices* **4** (1993), 93-106.
- [35] G. Segal. Categories and cohomology theories. *Topology* **13** (1974), 293-312.
- [36] C. Simpson. Higgs bundles and local systems. *Publ. Math. I.H.E.S.* **75** (1992), 5-95.
- [37] C. Simpson. Moduli of representations of the fundamental group of a smooth projective variety, I and II. *Publ. Math. I.H.E.S.* **79** and **80** (1994), 47-129 and 5-79.
- [38] C. Simpson. Flexible sheaves. Preprint q-alg 9608025.
- [39] C. Simpson. Homotopy over the complex numbers and generalized de Rham cohomology. *Moduli of Vector Bundles*, M. Maruyama (Ed.) *Lecture Notes in Pure and Applied Math.* **179**, Marcel Dekker (1996), 229-263.
- [40] C. Simpson. The Hodge filtration on nonabelian cohomology. Preprint alg-geom 9604005. To appear, *Proceedings of AMS summer conference, Santa Cruz*.
- [41] C. Simpson. A relative notion of algebraic Lie group and application to n -stacks. Preprint alg-geom 9607002.
- [42] C. Simpson. Algebraic (geometric) n -stacks. Preprint alg-geom 9609014.
- [43] C. Simpson. A closed model structure for n -categories, internal Hom , n -stacks and generalized Seifert-Van Kampen. Preprint alg-geom 9704006.
- [44] C. Simpson. Limits in n -categories. Preprint alg-geom 9708010.
- [45] C. Simpson. Effective generalized Seifert-Van Kampen. Preprint q-alg 9710011.
- [46] Z. Tamsamani. Sur des notions de n -catégorie et n -groupoïde non-strictes via des ensembles multi-simpliciaux. Thesis, Université Paul Sabatier, Toulouse (1996) available on alg-geom (95-12 and 96-07).

- [47] R. Thomason. Algebraic K-theory and etale cohomology, *Ann. Sci. E.N.S.* **18** (1985), 437-552.
- [48] C. Voisin. Variations de structure de Hodge et zéro-cycles sur les surfaces générales. Preprint, I.H.E.S., Mai 1993.